



Author Index

- Aagaard, P., 151
Abbasi, F., 1323
Abdelrahman, E., 91
Abudula, R., 1181, 1329
Adamkova, V., 1266, 1486
Adeli, K., 403, 689
Agardh, C.-D., 188
Agardh, E., 188
Agger, A., 1425
Agrawal, K.C., 645
Ahn, C.W., 206
Aigner, R., 862
Akahane, M., 995
Akahane, S., 995
Alacacioglu, A., 677
Ala-Rämi, A., 410
Alaupovic, P., 180
Albareda, M., 1115
Alemanno, N., 1474
Aliberti, G., 271
Allen, R., 351
Ambrosio, M.R., 1174
Amm-Azar, M., 947
An, C.-S., 1282
Anan, F., 552
Andersen, L.L., 151
Andersen, L.T., 1048
Andersen, O., 171
Andersen, U.B., 171
Andler, W., 397, 1155
Andres, R., 542
Angelini, V., 306
Angelopoulos, N., 1202
Ao, Y., 1480
Araki, A., 809
Araki, T., 381
Archer, W.R., 1390
Ardigò, D., 1420
Aregawi, D., 1345
Arii, K., 142
Arita, K., 699
Arnaud-Battandier, F., 1108
Artieda, M., 1230
Asakawa, H., 67
Askari, H., 876
Aso, Y., 345, 1225
Assis-Pandochi, A.I.d., 300
Athyros, V.G., 1065
Atlan, C., 1442
Attali, J.-R., 1290
Attia, S., 1290
Aybay, C., 1384
Azar, R.R., 947
Azuine, M.A., 1056
Azuma, J., 55
Azzolini, A.E.C.S., 300
Babaya, N., 529
Bachelard, H., 1454
Badeau, M., 1454
Badell, G., 1115
Bae, K.T., 561
Baek, K.H., 49
Bahi, L., 634
Baik, S.H., 1510
Baker, L., 91
Balasubramanyam, M., 1002
Ballas, C., 1243
Bang, P., 275
Baquero, M., 33
Barakat, H., 1368
Barakat, H.A., 1542
Bard, M., 1337
Barmada, M.M., 1552
Barré, P.E., 835
Barrett, P.H.R., 361
Bärtsch, P., 815
Batta, A.K., 38
Baynard, T., 866, 989
Beaudeau, J.-L., 829
Behre, C.J., 439
Beil, M., 522
Beliard, S., 1442
Bendikssen, A., 748
Benedict, C., 1356
Benzi, L., 1541
Bergamaschi, A., 891
Bergeron, J., 1014
Bergeron, J., 1390
Bergeron, N., 1390
Bergman, R.N., 902
Bergmann, K.v., 335
Berlan, M., 122
Bernard, E.A., 515
Bernardini, I., 719
Berneis, K., 227
Bertacca, A., 1541
Berthiaume, N., 735, 1515
Bhanot, S., 848
Biarnés, J., 235
Bierhaus, A., 815
Bigard, A.-X., 634
Billensteiner, E., 935
Birtles, T., 610
Bizeau, M.E., 1189
Blackett, P.R., 180
Blackman, M.R., 1162
Blanco-Vaca, F., 560
Blendea, M.C., 1337
Blevins, K.S., 180
Blouin, K., 1034
Blümer, R.M.E., 60
Bobkova, D., 1266
Boden, G., 1142
Boger, R., 351
Bollineni, J.S., 38
Bonadonna, S., 1174
Bondanelli, M., 1174
Bonfont-Rousselot, D., 829
Borkenstein, M., 862
Born, J., 391, 1356
Bornet, H., 584
Boston, R.C., 902
Bottini, E., 891
Bottini, N., 891
Bouchard, C., 1034
Bourgoin, F., 1454
Bourgouin, P.E., 1362
Brandauer, J., 97

- Bray, G.A., 24, 1506
 Bray, M.S., 910
 Briody, J., 500
 Brismar, K., 275
 Brun, J.-F., 1102, 1411
 Bryhni, B., 748
 Buckley, A.J., 500
 Bukowska, H., 72
 Burge, M.R., 460

 Caballero, A., 1115
 Caballero, B., 1162
 Cabero, L., 1504
 Caliskan, S., 677
 Calvo, L., 1230
 Cameron-Smith, D., 952
 Camino, N., 33
 Campbell, L.V., 821
 Campbell, W.W., 769
 Carey, M.F., 1269
 Carhart, R., 866
 Carhart, R. Jr., 989
 Carlson, C.J., 895, 1515
 Casals, E., 1504
 Casaschi, A., 403
 Casella, G.A., 841
 Caso, G., 841
 Cassader, M., 1474
 Caumo, A., 79
 Cavallo, M.G., 930
 Cecchetti, P., 1541
 Ceci, F., 271
 Cenarro, A., 1230
 Cendros, J.M., 1276
 Ceylan, C., 677
 Cha, B.S., 206
 Chambers, S.T., 1
 Champagne, C., 1506
 Chan, J., 1075
 Chang, Y., 1498
 Chanussot, F., 1087
 Chaparro, J., 561
 Chapot, R., 634
 Charlesworth, J.A., 821
 Chattergoon, N.N., 645
 Cheltowski, K., 72
 Chen, C.-C., 471
 Chen, J., 1329
 Chen, W.Y., 445
 Chen, X., 848
 Cheng, W.-T., 157
 Chevalier, S., 1529
 Chiba, H., 1536
 Chiellini, C., 1541
 Chien, Y.-H., 1467
 Chiu, C.-Y., 1082

 Chiu, Y.T., 445
 Cho, Y.M., 314
 Choi, D.S., 1510
 Choi, K.M., 1510
 Choi, M.G., 49
 Choi, Y.-H., 1448
 Christon, R., 1454
 Chrousos, G.P., 1350, 1472
 Chu, C.-H., 157
 Chung, M., 1411
 Chung, M.-Y., 1282
 Chusho, H., 1490
 Ciaccia, A.V., 939
 Ciccarone, A., 1541
 Cittadini, A., 306
 Civeira, F., 1230
 Cizza, G., 918
 Clarenbach, J.J., 335
 Clark, P.A., 1374
 Clore, J.N., 756
 Cobelli, C., 79
 Cochran, E., 255
 Coeli, C.M., 321
 Colleran, K.M., 460
 Colombo, M., 1181, 1329, 1425
 Comlekci, A., 677
 Conceição, F.L., 321
 Connelly, P.W., 103
 Coppolino, G., 930
 Corcoy, R., 1115
 Costanza, F., 930
 Couillard, C., 856, 1034
 Couture, P., 856, 1014, 1390
 Cozzolino, D., 306
 Cramer, R.M., 151
 Crispi, F., 1504
 Csako, G., 918
 Cubero, J.M., 560
 Cunnane, S.C., 1127
 Cupples, L.A., 1435
 Cynober, L., 1108

 da Silva, E.M.C., 321
 Dagogo-Jack, S., 876
 D'Agostino, R.B. Jr., 264
 Daimon, M., 669
 Daly, M.A., 1480
 Damberg, G., 1008
 Damcott, C.M., 1552
 Danielsson, A., 781
 Davidson, M.B., 1408
 Davidson, M.H., 939
 de Jonge, L., 24, 1506
 de Leiva, A., 1115
 de Lima Oliveira Brasil, R.R., 321
 Deacon, R., 522

 Debrececi, B., 1436
 DeCarlo, S.O., 848
 Deepa, R., 476
 degli Uberti, E.C., 1174
 Deiter, G., 212
 Del Prato, S., 1541
 Dela, F., 1398
 DeLany, J.P., 1506
 Delgado, J., 1504
 Dellow, W.J., 1
 Delsignore, R., 1420
 deLuca, C., 361
 Demarest, K., 848
 Denkins, Y., 1506
 DePaoli, A.M., 255
 Dériaz, O., 1390
 DeRiso, L., 1480
 Després, J.-P., 1034
 Desroches, S., 1390
 Di Giovanni, C., 271
 Didangelos, T.P., 1065
 Diehl, C.J., 1513
 Diez, J.G., 645
 Dimcevski, G., 165
 Djahangirian, O., 1362
 Döbrönte, R., 579
 Dodt, C., 1356
 Doga, M., 1174
 Doğan, Y., 800
 Dolnikowski, G.G., 361
 Domingo, N., 1087
 Domínguez, C., 1504
 Doshi, S., 351
 Dotti, M.T., 335
 Draznin, B., 1490
 Drinkard, B.E., 918
 Dunne, J.B., 645
 Dyrskog, S.E.U., 1181, 1316, 1329

 Ebihara, K., 1490
 Ebine, N., 508
 Eguchi, T., 370
 Ehrenkranz, J., 1407
 Eid, J., 1362
 El Matary, W., 15
 Elisaf, M., 1065
 Elisaf, M.S., 559
 Emoto, M., 330, 381
 Endert, E., 60
 Engelhardt, M., 542
 Erlich, J.H., 821
 Eskandari, F., 918
 Espinoza, J., 1027
 Eydoux, N., 634
 Fagerberg, B., 139, 439
 Fain, J.N., 220, 1546

- Fairow, C., 508
 Fan, J., 132
 Fang, C., 1142
 Farin, H.M. F., 1323
 Farkas, V., 1436
 Farnier, C., 1087
 Farthing-Nayak, P.J., 188
 Faulkner, D., 103
 Fazio, S., 1309
 Federico, A., 335
 Fédou, C., 1102
 Fehm, H.L., 391, 1356
 Feingold, E., 1552
 Felix, B., 227
 Fernandez, M.L., 1133
 Fernández-Castañer, M., 235
 Fernández-Real, J.M., 235
 Fernandez-Real, J.-M., 584
 Fernhall, B., 866
 Ferreira, L.D.M.C.-B., 1420
 Ferrell, R.E., 1552
 Figueroa, A., 866
 Finley, K.B., 902
 Fioriti, E., 930
 Fischer, G.M., 1436
 Florys, B., 1020
 Fogelstrand, L., 139
 Folch, N., 1428
 Fong, I., 103
 Fonseca, N.A., 645
 Fonseca, V.A., 645
 Fontaine-Bisson, B., 1390
 Fóris, G., 1147
 Fornage, M., 910
 Fournier, P.A., 1420
 Fox, C.S., 1435
 Franklin, R.M., 989
 Franzini, L., 1420
 Fredrikson, G.N., 1267
 French, D.N., 657
 Friedman, M.I., 421
 Frölich, M., 705
 Fujii, M., 775
 Fujimoto, M., 1490
 Fujimoto, S., 1469
 Fujisawa, T., 387, 529, 628
 Fujita, K., 85
 Fujita, M., 85
 Fujiwara, S., 381
 Fukui, T., 1447
 Fukumoto, S., 330
 Fukushima, M., 376, 1297, 1447
 Futenma, A., 1441
 Gagné, C., 1014
 Gagné, J.-M., 1014
 Gahl, W.A., 719
 Galbo, H., 1425
 Galitzky, J., 122
 Gallistl, S., 862
 Galvin, V.B., 1368
 Gambino, R., 1474
 Ganley, K.J., 793
 Gannagé-Yared, M.-H., 947
 Gao, J., 522
 Garcia, M.d.M., 235
 García-Iglesias, M.F., 741
 Garige, M., 729
 Garlick, B.A., 841
 Garlick, P.J., 841
 Gelain, D.P., 515
 Gennings, C., 756
 Genovese, D.J., 902
 George, P.M., 1
 Gepp, H., 881
 Gerber, L.H., 719
 Gerique, J.G., 33
 Germanos-Haddad, M., 947
 Geyer, G., 935
 Ghafoorunissa, 240
 Ghanaat, F., 604
 Ghiu, I.A., 97
 Giacca, A., 1250
 Giannopoulou, I., 866
 Giustina, A., 1174
 Gloria-Bottini, F., 891
 Glowinska, B., 1020
 Glueck, C.J., 113, 1345
 Goff, D.C. Jr., 264
 Goicoechea, J., 1230
 Gola, M., 1174
 Gold, P.W., 918
 Golden, M., 835
 Goldenberg, N., 113
 Goldfine, I.D., 598
 Gómez, A., 657
 Gong, M., 729
 Goodfellow, J., 351
 Gordon, P., 255
 Gorini, A., 983
 Gougeon, R., 1529
 Goula, A., 1202
 Gouloupoulou, S., 989
 Gounarides, J., 522
 Goya, K., 85
 Gragnoli, C., 786, 983
 Gratacós, E., 1504
 Gregersen, S., 1316, 1425
 Grès, S., 1087
 Grill, V., 247
 Grosclaude, J., 1087
 Gross, M.D., 910
 Gruetter, R., 1008
 Gu, J., 1453
 Guallar, A., 1230
 Guan, Y., 590
 Guardasole, V., 306
 Gündüz, M., 1384
 Guo, C.-Y., 1435
 Guo, Q., 689
 Gutterman, D., 764
 Habener, J.F., 786, 983
 Haffner, S.M., 264
 Hagberg, J.M., 97
 Halaby, G., 947
 Hall, K., 275
 Hallschmid, M., 1356
 Hama, S., 699
 Hamada, T., 430
 Hamajima, N., 1302
 Hamidi, M., 103
 Hamman, R.F., 1552
 Hanefeld, M., 1168
 Hansen, B.R., 171
 Haque, M., 1345
 Harding, L.K., 610
 Harman, S.M., 1162
 Harris, C., 1309
 Hasanoğlu, A., 1384
 Hashimoto, K., 142
 Hassinen, I.E., 410
 Hasty, A.H., 1309
 Hatsuda, S., 330, 381
 Hattori, S., 482
 Hattori, Y., 482
 Haub, M.D., 769
 Hauffen, J., 881
 Haugaard, S.B., 171
 Hayakawa, K., 995
 Hayashi, M., 370
 Hayashi, T., 1490
 Helge, J.W., 1048
 Hellström-Lindahl, E., 247
 Henkel, E., 1168
 Hennig, R., 881
 Henriksen, E.J., 1513
 Hergenç, G., 800
 Herman, R.M., 793, 1162, 1181, 1316, 1329, 1425
 Herron, K., 1133
 Hickner, R.C., 960, 1368, 1542
 Higa, M., 788
 Higaki, Y., 132
 Hirano, K.-i., 1095
 Hirao, K., 370
 Hiraoka, H., 1095
 Hirata, A., 669

- Hirose, H., 296, 370
Hitoglou-Makedou, A.D., 1236
Hoffer, L.J., 835, 1362
Holloszy, J.O., 960
Holm, J., 165
Holst, J.J., 1398
Hong, J., 1329
Home, M., 918
Hosoda, K., 1490
Hosoi, T., 809
Hosokawa, M., 1297
Houlden, H., 200
Houmard, J.A., 598
Hsia, S.H., 1408
Hsieh, P.-S., 157
Huang, A., 568
Huang, D., 568
Huang, J.-B., 1027
Huang, X., 1162
Huang, Y., 453
Hubacek, J.A., 1266, 1486
Hultén, L.M., 439
Hulthe, J., 139, 439
Hutten, B.A., 1499

Iannaccone, U., 891
Ibrahim, A., 240
Igarashi, M., 669
Ikeda, Y., 142
Ikegami, H., 387, 529, 628
Ikoma, A., 1461
Imura, I., 699
Inaba, M., 55
Inagaki, N., 1297, 1447
Inokuchi, T., 788
Inoue, K., 387, 529
Inoue, M., 281
Insull, W. Jr., 939
Inukai, T., 1225
Inukai, Y., 1225
Irani, N., 103
Isaji, M., 995
Ishida, K., 713
Ishihara, K., 775
Ishikawa, S.-e., 653, 1461
Ishimi, Y., 1536
Ishimura, E., 330
Ito, H., 809
Ito, Y., 1302
Itoi-Babaya, M., 529
Iversen, J., 171
Iwata, K., 1441

Jacobs, D.R. Jr., 910
Jacobson, L., 1259
Jaeger, R., 508

Jap, T.-S., 1082
Javor, E., 255
Jazet, I.M., 705, 1523
Jeanneret, C., 227
Jenkins, D.J.A., 103
Jenner, J.L., 361
Jenq, S.-F., 1082
Jensen, C.B., 1398
Jenssen, T.G., 748
Jentjens, R.L.P. G., 610
Jeppesen, P.B., 1181, 1329, 1425
Jeter, J.R. Jr., 645
Jeukendrup, A.E., 610
Ji, H., 421
Jia, D., 619
Jia, X., 508
Jimbu, Y., 669
Jin, N., 590
Jones, J.E., 821
Jones, P.J.H., 508
Jordan, H., 1411
Joseph, T., 1533
Judelson, D.A., 657
Jung, H.S., 314
Jung, J.-H., 1448
Jurva, J., 764

Kadowitz, P.J., 645
Kager, P.A., 60
Kaiser, G.R.R.F., 515
Kaiser-Kupfer, M.I., 719
Kajdaniuk, D., 1464
Kaji, Y., 775
Kakafika, A.I., 1065
Kameda, W., 669
Kamei, N., 194, 713
Kanaki, T., 573
Kanaley, J.A., 866
Kanaley, J.A., 989
Kang, E.S., 206
Kang, J.-S., 1282
Kang, M.I., 49
Kang, S.-H., 1282
Kangasmaa, H., 548
Kanno, Y., 370
Karagiannis, A., 1065
Kasai, K., 482
Kasayama, S., 55, 85
Kastelein, J.J. P., 1499
Kato, T., 669
Kavouras, S.A., 1472
Kawabata, Y., 387, 529
Kawabata, Y., 628
Kawabe, H., 370
Kawaguchi, A., 466
Kawaguchi, Y., 628

Kawakami, M., 1461, 653
Kawamitsu, H., 775
Kawamori, R., 1250
Kawano, T., 1461, 653
Kawasaki, Y., 1297
Kawase, I., 55, 85
Kawase, I., 85
Kazakos, K.A., 1236
Kazumi, T., 466
Keech, C.A., 939
Keizer, H., 1523
Keleş, İ., 800
Kelleher, J.K., 335
Kemeny, G., 1368
Kendall, C.W.C., 103
Kerbl, R., 862
Kern, W., 391, 1356
Keserü, B., 500
Khalifé, S., 947
Kiess, W., 397
Kim, D.-J., 1448
Kim, H.Y., 1510
Kim, H.J., 206
Kim, J.S., 1513
Kim, K.R., 206
Kim, K.-W., 1448
Kim, N.H., 1510
Kim, S.G., 1510
Kim, S.Y., 314
Kim, T., 1250
Kim, Y.S., 925
Kimoto, E., 330
Kimura, M., 669
Kimura, T., 430
Kissler, H.J., 881
Kissoon, N., 492
Kitajima, S., 132
Kitaoka, T., 699
Kjær, M., 151
Klein, R.L., 453
Klein, S., 561
Klement, P., 1250
Kleta, D., 1065
Knuuttila, M.L.E., 548
Kobayashi, K., 1469
Kobayashi, N., 1490
Kobayashi, Y., 1469
Kobyliansky, E., 975
Koehler, C., 1168
Koga, M., 55
Kogure, A., 488
Kohen Avramoglu, R., 689
Kohno, N., 194, 713
Komuro, R., 1095
Koo, S.I., 1133
Kopplin, P., 103

- Korach-André, M., 522
 Korytkowski, M.T., 1480
 Koseki, M., 1095
 Kos-Kudla, B., 1464
 Kosztáczky, B., 1147
 Kotila, C., 918
 Koulmann, N., 634
 Kovács, P., 579
 Koyama, H., 330, 381
 Kraemer, W.J., 657
 Kraml, P.J., 453
 Krettek, A., 139
 Kriketos, A.D., 821
 Kristensen, L.L., 165
 Krug, E.I., 1480
 Kruhoefter, M., 1425
 Kuivenhoven, J.-A., 1499
 Kulaputana, O., 97
 Kulkarni, P.M., 939
 Kulp, J.L., 598
 Kumon, Y., 142
 Kurisu, K., 699
 Kuroe, A., 376, 1297
 Kurose, T., 1447
 Kusakabe, T., 1490
 Kushwaha, R.S., 1075
 Kusunoki, M., 1441
 Kwok, M.G.K., 188
- Lafontan, M., 122
 Laguna, F., 741
 Lahoz, C., 33, 741
 Lai, W.-A., 1524
 Lairon, D., 1442
 Lakshman, M.R., 729, 1056
 Lam, T.K.T., 1250
 Lamarche, B., 856, 1390
 Lamarche, M., 1529
 Lambert, K., 634
 Lamou-Fava, S., 361
 Landry, N., 1390
 Lanska, V., 1266
 Lánská, V., 1486
 Lasaridis, A.N., 1236
 Lathrop, D.A., 1411
 Lau, J., 1411
 Laurent, D., 522
 Laurenza, I., 1541
 Lavoie, C., 1428
 Le, A., 764
 Ledet, T., 287
 Lee, B.-W., 1448
 Lee, D.-H., 1282
 Lee, E.T., 180
 Lee, H.C., 206
 Lee, H.K., 314
- Lee, I.-T., 1467
 Lee, K.W., 1510
 Lee, K.-U., 925
 Lee, M.-K., 1448
 Lee, M.-S., 1448
 Lee, W.-J., 1467
 Lee, W.J., 445
 Lee, W.L., 1467
 Lee, W.-Y., 1498
 Lee, W.Y., 49
 Lee, Y.-K., 1282
 Lefevre, M., 1506
 Legrand, A., 829
 Leonhardt, W., 1168
 Lepiorz, M., 1356
 Lever, M., 1
 Levy, C., 918
 Levy, D., 1435
 Lewis, M., 351
 Lewitt, M.S., 275
 Li, C.-I., 471
 Li, E., 561
 Li, G., 568
 Li, H., 568
 Li, L., 1142
 Li, Q., 1142
 Li, T.-C., 471
 Li, X., 568
 Li, Y., 24
 Liang, Y., 848
 Lichtenstein, A.H., 361, 1411
 Lim, S.K., 206
 Lima, J.J., 492
 Lin, C.-C., 471
 Lin, J., 568
 Lin, S.-Y., 1524
 Lin, S.Y., 445
 Lin, T.M., 1467
 Lindenthal, B., 335
 Linton, M.F., 1309
 Liu, C.-S., 471
 Liu, E., 132
 Liu, H., 568
 Liu, J., 876
 Liu, P.-H., 1524
 Livshits, G., 975
 Llurba, E., 1504
 Lobo, P.M., 321
 Lofgren, I., 1133
 Loftspring, M., 113
 Loh, C.-H., 157
 Loï, C., 1108
 Lopes-Virella, M.F., 453
 López-Bermejo, A., 584
 Lorec, A.-M., 1442
 Lorec, A.M., 1087
- Lorneau, B., 1290
 Lovejoy, J.C., 1506
 Lucarelli, P., 891
 Lucisano-Valim, Y.M., 300
 Lynch, K., 188
- Maassen, J.A., 1523
 MacDonald, K.G., 598, 1368
 Madan, A.K., 220, 1546
 Maddux, B.A., 598
 Madsbad, S., 171, 1398
 Maeda, K., 775
 Maehata, E., 281
 Maeno, T., 381
 Maffei, M., 1541
 Magnusson, P., 151
 Magrini, A., 891
 Maiyoh, G.K., 403
 Manetta, J., 1411
 Manfrini, S., 930
 Mao, J., 590
 Marcell, T.J., 533
 Marcovina, S.M., 361
 Marek, B., 1464
 Maresh, C.M., 657
 Marette, A., 1454
 Marin, V., 1087
 Marino, G., 891
 Marliss, E.B., 1529
 Marshall, J.A., 1552
 Martinez, P.E., 918
 Martín-Gallán, P., 1504
 Maruyama, T., 370
 Massicotte, D., 1428
 Mastellone, I., 1087
 Masuzaki, H., 1490
 Matarazzo, M., 306
 Matsumoto, S., 1225
 Matsutomo, R., 1225
 Matsuura, F., 1095
 Matsuyama, A., 1095
 Matsuzawa, A., 995
 Matthán, N.R., 1411
 Mattila, P.T., 548
 Mauras, N., 492
 McAuley, K.A., 533
 McDowell, I., 351
 McNamara, D.B., 645
 Meguro, S., 67
 Meinders, A.E., 705, 1523
 Melançon, S., 1454
 Menschikowski, M., 1168
 Mercadé, I., 1504
 Mercier, J., 634, 1102, 1411
 Meriño-Ibarra, E., 1230
 Metz, L., 1102

- Mezey, G., 579
Mikhailidis, D.P., 1065
Miki, T., 330
Milionis, H.J., 559
Millar, J.S., 361
Millo, B., 72
Milord, E., 786
Min, Y.-K., 1448
Miserez, A.R., 227
Mistry, S., 918
Miyake, K., 775
Miyahara, F., 1490
Miyashita, K., 296
Miyata, T., 1441
Miyatake, A., 85
Miyazawa, T., 1490
Mizutani, K., 1441
Moat, S., 351
Modi, P., 930
Moffett, S.P., 1552
Mohan, A., 476
Mohan, V., 476, 1002
Mondal, D., 645
Monia, B.P., 848
Monteiro, S.C., 515
Monti, L., 1420
Montvilas, P., 165
Morais, J.A., 1529
Mori, T., 370
Morimoto, M., 132
Morimoto, Y., 85
Moritani, T., 430
Moro, C., 122
Most, M.M., 1506
Mostaza, J.M., 33, 741
Mouritsen, E.A., 165
Mouslech, T.F., 1236
Mozaffari, M.S., 1243
Muller, D.C., 542
Muntzel, M.S., 1533
Murthy, S.N., 645
Muser, J., 227
Musso, C., 255

Nadler, J.L., 1453
Nagai, N., 430
Nagai, Y., 788
Nagasaka, S., 1297
Naim, A.C., 1453
Naito, M., 1302
Naitoh, T., 573
Nakagawa, M., 552
Nakai, Y., 376, 1297
Nakamura, T., 653, 1095, 1441
Nakanishi, K., 788
Nakanishi, S., 194, 713

Nakao, K., 1490
Nakib, S., 1108
Namai, K., 653, 1461
Napoli, R., 306
Nareika, A., 453
Naruszewicz, M., 72
Nasr, I., 15
Nassis, G.P., 1472
Natarajan, S., 240
Nawroth, P.P., 815
Nayak, R.C., 188
Nebert, D.W., 568
Neemallah, R., 947
Németh, J., 579
Neufer, P.D., 1048
Neveux, N., 1108
Newcombe, R., 351
Niedziolka, D., 1464
Nielsen, J.O., 171
Nierman, M.C., 1499
Niessner, A., 935
Nilsson, J., 1267
Nilsson, P.M., 1236
Nishida, M., 1095
Nishimura, K., 55
Nishino, M., 529, 628
Nishio, K., 1302
Nishizawa, Y., 330, 381
Nogalska, A., 1041
Noh, J.-H., 1448
Nojima, H., 194
Nojima, K., 529
Nomura, T., 1441
Nowak, M., 1464
Numeroso, F., 1420
Nystrom, F.H., 781

Oana, F., 995
Obach, R., 1276
Obregon, D.F., 645
O'Brien, K., 719
Ogata, T., 370
Ogawa, H., 1441
Ogawa, T., 296
Ogawa, Y., 1490
Ogihara, T., 387, 529, 628
Oh, E.S., 49
Oh, K.W., 49
Oh, Y.-S., 1282
Ohama, T., 1095
Ohara, T., 775
Ohashi, A., 1469
Ohgushi, M., 376
Ohno, Y., 370
Ohnuma, H., 669
Ohtaki, M., 713

Ohya, M., 376, 1297
Oizumi, T., 669
Okamoto, S., 1447
Okubo, M., 194, 713
Okumura, K.-i., 345
Okuyama, A., 55
Olafsen, K., 748
Onat, A., 800
Onderdonk, A., 103
Ono, M., 529, 628
Onofri, A., 1174
Onuma, S., 788
Onwumere, O., 1533
Ooie, T., 552
Oppenorth, T.J., 735
Ordóñez-Llanos, J., 1115
Orimo, H., 809
Orntoft, T., 1425
Osada, T., 1048
Osaki, F., 142
Osborne, M.C., 848
Ota, K., 142
Otani, Y., 55
Otsuki, M., 619
Otvos, J.D., 264
Ou, H.-C., 1467
Ouslimani, N., 829
Outterbridge, L.V., 421
Ouwens, D.M., 1523
Ozanne, S.E., 500
Özbay, F., 1384

Pabst, K.M., 1162
Pacák, K., 1259
Paganelli, F., 1442
Pagano, G., 1474
Pagliassotti, M.J., 1189
Páll, D., 1147
Palmer, T.N., 1420
Pantsulaia, I., 975
Papageorgiou, A.A., 1065
Papantakou, K., 1472
Papassotiropoulos, I., 1350
Paragh, G., 579, 1147
Paragh, G. Jr., 1147
Pariès, J., 1290
Park, C.Y., 49, 925
Park, D.-B., 1282
Park, H.J., 314
Park, H.S., 925
Park, K.S., 314
Park, S.W., 49, 206
Parker, T., 103
Patel, C., 1243
Patel, S.J., 1056
Patterson, B., 682

- Peake, P.W., 821
 Péan, M., 1428
 Peczynska, J., 1020
 Peinnequin, A., 634
 Peitl, B., 579
 Peletidou, A., 1065
 Pelikanova, T., 122
 Peña, R., 33, 741
 Pender, C., 598
 Peng, X., 1453
 Peraire, C., 1276
 Pereira, R.I., 1490
 Pérez, A., 560
 Péronnet, F., 1428
 Perry, M.B., 719
 Perry, M.L.S., 515
 Peters, A., 391
 Petty, H.R., 1027
 Peynet, J., 829
 Phares, D.A., 97
 Piatti, P.M., 1420
 Pichler, G., 862
 Pijl, H., 705, 1523
 Pilegaard, H., 1048
 Pinach, S., 1474
 Pintó, X., 741
 Ploug, T., 1425
 Poehlman, E.T., 1267
 Polak, J., 122
 Poledne, R., 1486
 Polizello, A.C.M., 300
 Pomerleau, S., 856
 Poretsky, L., 200
 Pories, W.J., 1368
 Portugal, H., 1442
 Potteiger, J.A., 756
 Powderly, W.G., 561
 Pozzilli, P., 930
 Pradeepa, R., 476
 Pranikoff, J., 1345
 Prefaut, C., 1411
 Premanand, C., 1002
 Privette, J.D., 1542
 Proietta, M., 271
 Prud'homme, D., 1034
 Pugeat, M., 584
 Pulignano, I., 271
 Py, G., 634, 1102

 Quilliot, D., 966
 Quintana, E., 180
 Quirk, E.K., 561

 R. Seaquist, E., 1008
 Racette, S.B., 960
 Radha, V., 476

 Raghavan, S., 91
 Rao, A.V., 103
 Rao, M.N., 729
 Rasmussen, I.H., 165
 Rasmussen, L.M., 287
 Ratamess, N.A., 657
 Raynaud, E., 1102
 Reaven, G.M., 1323, 1420
 Reaven, P.D., 533
 Reeds, D.N., 561
 Reinehr, T., 397, 1155
 Rema, M., 1002
 Reynolds, J.C., 918
 Rhee, E.-J., 49, 1498
 Ricart, W., 235, 584
 Richter, B., 935
 Richterova, B., 122
 Rip, J., 1499
 Rizk, M.M., 15
 Rodriguez, A., 542
 Rodriguez, R., 1075
 Rodríguez-Espinosa, J., 1115
 Roemmich, J.N., 1374
 Rogol, A.D., 1374
 Romero, L.A., 460
 Romero, R., 1027
 Romijn, J.A., 705
 Rondinone, C.M., 895, 1515
 Rood, J., 1506
 Rosenthal, M.J., 1480
 Roth, C.L., 1155
 Rubin, B.I., 719
 Ruel, G., 856
 Ruiten, A.F.C., 60
 Ruiu, G., 1474
 Russo-Carbolante, E.M.d.S., 300
 Ryan, M.A.A., 1127
 Ryu, S., 1498

 Saboohi, F., 835, 1362
 Saccà, L., 306
 Saengsirisuwan, V., 1513
 Saheki, T., 1469
 Saikawa, T., 552
 Saito, I., 296, 370
 Saito, M., 653
 Saito, T., 653, 1461
 Sakai, N., 1095, 1499
 Sakane, N., 430
 Saklamaz, A., 677
 Salehi, M., 200
 Salen, G., 38
 Salles, E., 321
 Saltin, B., 1048
 Sampathkumar, R., 1002
 Sanchez, H., 634

 Sandor, A., 1436
 Sansoy, V., 800
 Sarafidis, P.A., 1236
 Saruta, T., 296, 370
 Sasvary, D., 841
 Sauerwein, H.P., 60
 Schaart, G., 1523
 Schaefer, E.J., 361
 Schaffer, S.W., 1243
 Schiappoli, A., 271
 Schonfeld, G., 682
 Schreiner, P.J., 910
 Schulpis, K.H., 1350
 Schultes, B., 391
 Schultz, K., 287
 Schuille, P.O., 881
 Scott, R.S., 1
 Sebring, N., 719, 918
 Seino, Y., 376, 1297, 1447
 Seman, L.J., 361
 Semenkovich, C.F., 561
 Sengenès, C., 122
 Seno, H., 699
 Seo, J.A., 1510
 Seres, I., 1147
 Shachter, N.S., 1133
 Shannon, K.A., 756
 Shannon, R.M., 756
 Sharman, M.J., 657, 1133
 Shaw, C., 610
 She, P., 848
 Shefer, S., 38
 Sheikh, A., 200
 Sherman, A., 113
 Sheu, W.H.-H., 445, 1467, 1524
 Shi, J., 568
 Shigenobu, M., 699
 Shih, K.-C., 157
 Shimbo, T., 1447
 Shin, C.S., 314
 Shinohara, K., 330, 381
 Shintani, M., 628
 Shoji, T., 330, 381
 Shuldiner, A.R., 97
 Siddhanti, S., 939
 Sidossis, L.S., 1472
 Sieja, K., 72
 Sieminska, L., 1464
 Silvestre, R., 657
 Singhal, P., 79
 Sirvent, P., 1102
 Sjöberg, S., 139
 Skenderi, K., 1472
 Skodova, Z., 1266, 1486
 Slany, J., 935
 Sloniger, J.A., 1513

- Slyper, A., 764
 Smith, S.R., 24, 1506
 Snow, R.J., 1269
 Soares, D.V., 321
 Soler, J., 235
 Soliman, A.T., 15
 Song, A.M., 1480
 Song, M.K., 1480
 Song, Y.D., 206
 Souza, L.F., 515
 Sowers, J.R., 1337
 Spiering, B.A., 657
 Spina, L.D.C., 321
 Stabler, S., 351
 Stafylas, P.C., 1236
 Stanojevic, V., 983
 Stanosz, S., 72
 Stathis, C.G., 1269
 Steffes, M.W., 910
 Stich, V., 122
 Stoddart, M., 180
 Stoffel-Wagner, B., 1155
 Storgaard, H., 1398
 Storlien, L.H., 1316
 Strålfors, P., 781
 Stroes, E.S.G., 1499
 Su, Z., 568
 Sucajtyś-Szulc, E., 1041
 Suciu, G., 165
 Suehiro, T., 142
 Suetta, C., 151
 Sugimura, K., 775
 Süheyl Ezgü, F., 1384
 Sumner, A.E., 902
 Sun, D., 522
 Sun, H., 132
 Sung, K.C., 1498
 Suwannarat, P., 719
 Suzuki, H., 1447
 Suzuki, K., 1302
 Suzuki, S., 281
 Swierczynski, J., 1041
 Swift, L.L., 1309
 Sylvester, J.E., 492
 Szilvassy, Z., 579

 Tabata, A., 1469
 Tabata, T., 330
 Taboada, M., 741
 Taguchi, M., 619
 Taha, A.Y., 1127
 Tahara, H., 330, 381
 Tai, Y.-H., 157
 Takaha, N., 55
 Takahashi, N., 552
 Takahashi, O., 1447

 Takakura, Y., 488
 Takaya, J., 1469
 Takebayashi, K., 1225
 Takeda, H., 995
 Taki, Y., 376, 1297
 Tamakoshi, A., 1302
 Tamemoto, H., 653, 1461
 Tanaka, T., 1490
 Tang, Y., 1142
 Tang, Y.-J., 1467
 Taniguchi, A., 376, 1297
 Taniguchi, H., 529
 Taniyama, M., 281
 Tanner, C.J., 598
 Tanoli, T., 682
 Tayek, J.A., 604
 Taylor, R., 79
 Tchernof, A., 1034
 Temiz, A., 677
 Thabet, A., 15
 Theriault, A.G., 403
 Thérond, P., 829
 Thomas, M.K., 983
 Thomas, W., 1008
 Thompson, C.H., 500
 Thompson, M., 500
 Tichansky, D.S., 1546
 Tkac, I., 1008
 Toda, H., 488
 Todd, M.K., 1218
 Tokimitsu, I., 67
 Tokunaga, K., 67
 Tolis, G., 1202
 Tominaga, A., 699
 Tominaga, M., 669
 Torella, R., 306
 Torvik, S., 918
 Tosini, F., 1087
 Tourkantonis, A.A., 1236
 Tracy, T., 1345
 Trappolini, M., 271
 Traustadóttir, T., 533
 Tremblay, A., 1034
 Tremblay, A.J., 1014
 Triandafillopoulou, M., 1472
 Tritapepe, L., 271
 Trocóniz, I.F., 1276
 Trofimov, S., 975
 Tsakiris, S., 1350
 Tsatsoulis, A., 559
 Tselepis, A.D., 559
 Tsilou, E., 719
 Tsujii, K.-i., 1095
 Tsuruzoe, N., 573
 Tsutsumi, K., 1441
 Tufekovic, G., 151

 Tümer, L., 1384
 Tunstall, R.J., 952
 Turcotte, L.P., 1218
 Türkmen, S., 800
 Tziomalos, K., 1065

 Uberti, B., 1474
 Uchino, H., 1250
 Umekawa, T., 488
 Upton, D.A., 460
 Urban, M., 1020
 Ushikai, M., 1469
 Utsumi, K., 1441

 Vaag, A., 1398
 Vaisman, M., 321
 Valensi, P., 1290
 Valero, R., 1442
 Valtueña, S., 1420
 van Raalte, D.H., 1499
 van Thien, H., 60
 VandeBerg, J.F., 1075
 VandeBerg, J.L., 1075
 Varga, Z., 1147
 Vary, T.C., 212
 Vega, G.L., 902
 Veldhuis, J.D., 1374
 Velmurugan, K., 476
 Verlaan, G., 151
 Vialettes, B., 1442
 Vianello, B., 1541
 Vice, E., 1542
 Vidgen, E., 103
 Vimalaswaran, K.S., 476
 vinh Thuan, D., 60
 Volaufova, J., 24
 Volek, J.S., 657, 1133
 Volund, A., 171
 von Eyben, F.E., 165
 von Eyben, R., 165
 Von Preussenthal, G.M., 983
 Vounatsou, M., 1350

 Wakabayashi, S., 1225
 Wang, C., 568
 Wang, G., 590
 Wang, H.-J., 471
 Wang, J., 492
 Wang, P., 113, 1345
 Wang, W., 180
 Wang, X., 590, 1536
 Wang, Y., 508
 Wang, Y.-Y., 1524
 Waring, R.H., 610
 Warren, B.J., 756
 Watanabe, A., 699

- Watanabe, H., 67
 Watanabe, T., 132
 Watson, G., 657
 Watts, L.M., 848
 Wei, J., 590
 Weidong, Y., 1441
 Weinrich, M., 91
 Weinstock, R.S., 866
 Weiss, C., 815
 Weiss, E.P., 97, 960
 Wells, A.M., 769
 Weltman, A., 1374
 Welty, F.K., 361
 Wen, W., 561
 Wessale, J.L., 735
 West, K., 1133
 Westergren, H., 781
 Weverling, G.-J., 60
 Wilhelm, O., 682
 Williams, W., 1250
 Willis, W.T., 793
 Wilson, J.W., 1480
 Wilson, P.W.F., 1435
 Wilttrout, S.A., 492
 Winer, N., 1337
 Winters, S.J., 91, 1480
 Wirth, S., 935
 Wohn, C.R., 97
 Wojciechowska, C., 1464
 Wolever, T.M.S., 103
 Woloszczuk, W., 935
 Wu, H.S., 445
 Wu, J., 1536
 Wu, Y.-C., 1082

 Xiao, C., 568
 Xiao, J., 1329, 1425

 Xie, H., 24
 Xu, D., 1420
 Xu, G., 38
 Xueping, E., 561

 Yablonskiy, D., 682
 Yagi, N., 67
 Yamada, K., 1536
 Yamada, N., 132
 Yamada, Y., 1447
 Yamaguchi, D.T., 1480
 Yamaguchi, H., 669
 Yamamoto, H., 85
 Yamamoto, K., 67, 1302
 Yamane, K., 194, 713
 Yamano, G., 376
 Yamashita, S., 1095
 Yamato, F., 1469
 Yanagawa, T., 376
 Yang, C.-H., 1467
 Yang, G., 1142
 Yang, H., 1480
 Yang, Q., 1435
 Yannakoulia, M., 1472
 Yano, M., 281
 Yarasheski, K.E., 561
 Yaspelkis, B.B. III, 1218
 Yasu, T., 653
 Yasue, S., 1490
 Yeshao, W., 1453
 Yesil, S., 677
 Ylihautala, M., 410
 Yokoyama, H., 381
 Yoo, H.J., 49
 Yoon, K.H., 49
 Yoshida, N., 345
 Yoshida, T., 488

 Yoshii, H., 1250
 Yoshii, S., 376, 1297
 Yoshikawa, H., 247
 Yoshikawa, T., 488
 Yoshimatsu, H., 552
 Yoshino, G., 466
 Yoshioka, K., 488
 Yotsumoto, T., 573
 Youn, B.-S., 314
 Young, J., 255
 Youngren, J.F., 598
 Yovos, J.G., 1236
 Yu, K.-Y., 314
 Yue, P., 682
 Yukioaka, H., 1490
 Yun, E.J., 49
 Yun, Y.S., 206

 Zannad, F., 966
 Zarra, E., 306
 Zatelli, M.C., 1174
 Zavaroni, I., 1420
 Zebis, M.K., 151
 Zern, T., 1133
 Zhang, L., 1142
 Zhang, S., 568
 Zhang, Y., 729
 Zheng, K., 568
 Zhou, B., 568
 Zhu, B., 1345
 Zhu, M.-y., 1309
 Ziegler, O., 966
 Ziegler, S., 935
 Zinker, B.A., 735, 1515
 Zochowska, E., 72
 Zotter, H., 862

Subject Index

- A30400, p38 inhibitor, effect on glucose uptake in insulin-resistant 3T3-L1 adipocytes, 895-901
- AA, *see* Amino acids
- Abdominal adipose tissue (AT), and serum levels of prothrombic factors in younger, premenopausal women with MDD, 918-924
- Abdominal symptoms, subjective, and postprandial hyperglycemia, effects of voglibose and acarbose, 387-390
- Acarbose, and voglibose, effects on postprandial hyperglycemia and subjective abdominal symptoms, 387-390
- ACE inhibitors, *see* Angiotensin-converting enzyme inhibitors
- Acetaldehyde, effect on 2,6-ST mRNA expression in wild-type HepG2 liver cells, 733
- Acidosis, effects on tissue protein synthesis in rats, 841-847
- Acid phosphatase locus 1 (ACP₁) genotype, relation to serum glucose concentration in healthy adult subjects, 891-894
- ACP₁, *see* Acid phosphatase locus 1
- Acute respiratory acidosis, effects on tissue protein synthesis in rats, 841-847
- S-Adenosylhomocysteine, relationship to S-adenosylmethionine, asymmetric dimethylarginine, and endothelial function, in healthy human subject during experimental hyper- and hypohomocysteinemia, 351-360
- S-Adenosylmethionine, relationship to S-adenosylhomocysteine, asymmetric dimethylarginine, and endothelial function, in healthy human subject during experimental hyper- and hypohomocysteinemia, 351-360
- Adipocyte plasma membrane fatty acid composition, and insulin sensitivity, effects of dietary *trans*-FA on, 240-246
- Adipocyte(s)
- effect of insulin on release of VEGF, IL-8, PAI-1, and IL-6 by, 220-226
 - effect of obesity on TGF β 1 release by human AT, 1546-1551
 - omental, *see* Omental adipocytes
 - subcutaneous, *see* Subcutaneous adipocytes
 - 3T3-L1
 - insulin-resistant, effect of p38 MAP kinase inhibition on glucose uptake in, 895-901
- Adipocyte(s) (continued)
- insulin-stimulated adiponectin secretion from, effect of inhibition of PI3K signaling pathway, 1636-1643
- Adipocytokines
- and inflammatory cytokine levels, effects of diet and/or exercise in postmenopausal NIDDM women, 866-875
 - levels, and S_i, effect of pravastatin in healthy nondiabetic patients, 947-951
 - plasma, *see* Plasma adipocytokines
 - reciprocal association with insulin resistance and C-reactive protein, in clinically healthy men, 439-444
- Adiponectin
- circulating, *see* Circulating adiponectin
 - or C-reactive protein, association with exercise training, 533-541
 - effect of pravastatin in healthy nondiabetic patients, 947-951
 - and glucose production, in patients infected with *Plasmodium falciparum*, 60-66
 - and insulin sensitivity in overweight and obese girls, effects of aerobic exercise training, 1472-1479
 - insulin-stimulated secretion from 3T3-L1 adipocytes, effect of inhibition of PI3K signaling pathway, 1636-1643
 - low levels of, association with metabolic syndrome, 476-481
 - physiologic difference between obese (fa/fa) and lean Zucker rats, 995-1001
 - plasma, *see* Plasma adiponectin
 - protective effect against oxidative stress in Japanese Americans, 194-199
 - reciprocal association with insulin resistance and C-reactive protein, in clinically healthy men, 439-444
 - role in neurophysiologic modulation of obesity, 1206-1207
 - serum, *see* Serum adiponectin
- Adiponectin-leptin ratio, correlation with parameters of insulin resistance, in NIDDM, 281-286
- Adiponectin mRNA (messenger ribonucleic acid), in chronic blockade of NO synthesis, 482-487
- Adiponectin receptor 1, mRNA expression in obese (fa/fa) and lean Zucker rats, 998

Adipose tissue (AT)

- abdominal, *see* Abdominal adipose tissue
- brown, *see* Brown adipose tissue
- depots, and food consumption and body weight, effects of camostat, 621-622
- eNOS content in, from obese and lean African American and white American women, 1368-1373
- gene expression in, and hyperglycemia prevention in Zucker diabetic fatty rats, effects of exercise training on, 1571-1581
- in obesity, *see* Adipose tissue, in obesity
- relationship to plasma adiponectin, in end-stage renal disease, 330-334
- subcutaneous, *see* Subcutaneous adipose tissue
- visceral, subcutaneous, and total, effect of TZDs on, 24-32
- white, *see* White adipose tissue
- see also* entries beginning with element: Adip-

Adipose tissue (AT), in obesity

- insulin sensitivity and, effects of aerobic exercise training, 1472-1479
- role in neurophysiologic modulation of obesity, 1203-1205
- TGF β 1 release by, 1546-1551

Adiposity

- excess, relation to IMGU, as determined by BMI and WC, 1323-1328
- and insulin resistance, influence on lipoprotein size and subclass concentrations, 264-270
- longitudinal measures of, inverse effects of PPAR γ 2 Pro12Ala polymorphism in African Americans and whites, 910-917

ADMA, *see* Asymmetric dimethylarginine**Adolescents**

- boys, with CDGP versus those with β -thalassemia major, hCG therapy in, 15-23
- Cherokee Indians, apo C-III bound to apo B-containing lipoproteins and insulin resistance in, 180-187
- and children, *see* Adolescents, children and
- obese, insulin sensitivity indices of glucose and free fatty acid metabolism, in relation to serum lipids in, 397-402

Adolescents, children and

- with obesity, hypertension and IDDM, levels of sICAM-1, sVCAM-1 and selectins, 1020-1026
- relationship between fasting plasma ghrelin and metabolic parameters in, 925-929

Adrenal insufficiency, diagnosis in a patient with Allgrove syndrome and ALADIN gene mutation, 200-205**Adrenal sympathetic nerve activity (SNA),**

- hypoglycemia-induced, effect of captopril on reflex increases in, 1679-1686

Adrenergic receptor, lipolytic pathways dependent on ANP

- and, differential regulation in human SAT, 122-131

 β_2 (beta $_2$) adrenergic receptor haplotype, and sex, effect on resting and terbutaline-stimulated whole body lipolysis, 492-499**Adults**

- with growth hormone deficiency after long-term growth hormone replacement, carotid artery IMT and lipid profile in, 321-329

- healthy subjects, relationship between serum glucose concentration and ACP $_1$ genotype in, 891-894

Advanced glycation end products (AGEs), *see* Serum advanced glycation end products**Advanced glycation index, association with NIDDM and microangiopathy, 1002-1007****Aerobes, antibiotics against, effects on serum lipids and colonic microflora, 103-112****Aerobic exercise training, effects on insulin sensitivity, body weight, body fat, adiponectin and inflammatory markers in overweight and obese girls, 1472-1479****African Americans, *see* Race and ethnicity****Age**

- association with circulating OPG in healthy Korean women, 49-54

- correlation to plasma leptin levels, in children with short stature administered with insulin, 862-865
- and declining androgen levels, contribution to features of MS in men, 1034-1040

- of intramyocellular lipids, role in progression of insulin resistance in nondiabetic Zucker rats, 522-528

- see also* specific age groups; for example: Adolescents, Children

Aged, healthy men and women, effects of GH and/or sex steroids on whole-body protein turnover, 1162-1167**AGEs, *see* Serum advanced glycation end products****Aging**

- disabilities associated with, S1:40

- effects on substrate oxidation during exercise in middle-aged and young athletes versus sedentary men, 1411-1419

- WAT in, effect of leptin on lipogenic enzyme expression through modification of SREBP-1c gene expression, 1041-1047

Akt phosphorylation, and glucose transport, effect of continually high insulin levels in human myoblasts, 1687-1693**ALA, *see* Alpha (α)-linolenic acid****ALADIN gene mutation, and Allgrove syndrome, diagnosis of adrenal insufficiency in a patient with, 200-205****Alanine aminotransferase (ALT)**

- normal concentrations, hyperinsulinemia predicts hepatic fat content in healthy individuals with, 1566-1570

- plasma, *see* Plasma alanine aminotransferase
- relation to serum ferritin and MS in Korean postmenopausal women, 1510-1514

Albuminuria

- and hypertension, association with circulating APAAs, in NIDDM, 188-193

- see also* Microalbuminuria

Alcohol

- mechanism of action in down-regulation of Gal β 1, 4GlcNAc α 2,6-sialyltransferase mRNA, in human liver cell lines, 729-734
- and xylitol, effect of simultaneous dietary administration on bone resorption, 548-551
- Alcohol administration, long-term, effect on synthesis of myofibrillar and sarcoplasmic proteins in heart, 212-219
- Alcohol drinking, and walking time for high serum uric acid levels, possible interactions with eNOS genotype among Japanese, 1302-1308
- Alcohol feeding, chronic, effects on activities and RSRs of various sialidases in rat liver, 1056-1064
- Alfacalcidol, versus alendronate, effects on BMD and bone metabolic markers in PM asthmatic women treated with inhaled corticosteroids, 85-90
- Alkaline phosphatase (ALP), *see* Plasma alkaline phosphatase
- Alkaptonuria, urinary HGA levels in, effect of nitisinone, 719-728
- Allgrove syndrome, and *ALADIN* gene mutation, diagnosis of adrenal insufficiency in a patient with, 200-205
- Allopurinol
 - effect on plasma adiponectin, in chronic blockade of NO synthesis, 482-487
 - influence on urinary purine loss after repeated sprint exercise in man, 1269-1275
- Alloxan, DM induced by, effects of FK-614 on peripheral glucose utilization and hepatic insulin extraction in dogs, 1250-1258
- ALP, *see* Plasma alkaline phosphatase
- Alpha (α)-glucosidase inhibitors, voglibose and acarbose, effects on postprandial hyperglycemia and subjective abdominal symptoms, 387-390
- 1-Alpha (α)-hydroxyvitamin D₃, versus alendronate, effects on BMD and bone metabolic markers in PM asthmatic women treated with inhaled corticosteroids, 85-90
- Alpha (α)-linolenic acid (ALA), systematic review and meta-analysis of impact on selected arrhythmia outcomes in animal models, 1557-1565
- Alpha (α)-lipoic acid (LA), supplementation with, effects on exercise-induced blood coagulation activation, 815-820
- Alpha (α)-melanocyte-stimulating hormone, role in neurophysiologic modulation of obesity, 1208
- Alpha,beta (α,β)-unsaturated aldehyde, 4-hydroxy-2-nonenal (HNE), effect on 2,6-ST mRNA expression in CYP2E1 cells, 733
- ALT, *see* Alanine aminotransferase
- Aminergic receptors, and glucocorticoids, neurobiological basis of depression, S1:26
- Amino acids (AA), metabolism by cultured rat Sertoli cells, 515-521
- p*-Aminomethylbenzoic acid (PAMBA), effect of aromatic ring-containing drugs on carnitine biosynthesis, 1582-1586
- Amphetamine transcript, role in neurophysiologic modulation of obesity, 1208-1209
- Anaerobes, antibiotics against, effects on serum lipids and colonic microflora, 103-112
- Androgen-deprivation therapy, for prostate cancer, hyperglycemia after, and effect of pioglitazone, 55-59
- Androgen(s)
 - declining levels, and age, contribution to features of MS in men, 1034-1040
 - excess, contribution to CV risk profile in postmenopausal women with NIDDM, 1626-1631
 - male mice deficient in, effect of *Puerariae radix* on bone loss, 1536-1541
- Angiotensin-converting enzyme (ACE) inhibitors, in targeting CV risk in nondiabetic patients with MS, 1065-1074
- Angiotensin II, oxidative burst induced by, sensitivity to fluvastatin in neutrophils of hypercholesterolemic patients, 1147-1154
- Angiotensin II AT1 receptor (AT1R), functional interaction with AT2R, in fructose-induced insulin resistance and hypertension, 157-164
- Angiotensin II AT2 receptor (AT2R), functional interaction with AT1R, in fructose-induced insulin resistance and hypertension, 157-164
- Angiotensin II receptor, effect of selective antagonism on whole-body insulin sensitivity and muscle glucose transport in hypertensive TG(mREN2)27 rats, 1659-1668
- Animal models, of IR, metabolic responses with ET antagonism in, 735-740
- Anionic peptide factor (APF), effects on UC efflux, 1087-1094
- Anorexia, hypoxia-induced, and RV hypertrophy, role in lactate transport and monocarboxylate transporter expression in rat muscle, 634-644
- ANP, *see* Atrial natriuretic peptide
- ANS, *see* Autonomic nervous system
- Anthropometric parameters, relationship to serum resistin level, among healthy subjects, 471-475
- Antibiotics, effect on serum lipids and colonic microflora, 103-112
- Antihypertensive drugs, in targeting CV risk in nondiabetic patients with MS, 1065-1074
- Antioxidant capacity, plasma, and OxLDL levels in men after short-term cranberry juice consumption, 856-861
- Antipericyte autoantibodies (APAAAs), circulating, association with albuminuria and hypertension in NIDDM, 188-193
- Antisense oligonucleotides (ASO), against GCCR, effects on hepatic glucose production and hyperglycemia in DM mice, 848-855

- Aortic endothelial cells (ECs), effect of MET on intracellular production of ROS in, 829-834
- APAAs, *see* Antipericyte autoantibodies
- APF, *see* Anionic peptide factor
- Apo(s), *see* specific apolipoproteins
- APOA* variant, (Val153 > Met), sex-specific effect on plasma HDL-C, 1632-1635
- Apolipoprotein (Apo), effects of raloxifene and low-dose simvastatin coadministration in PM women with primary hypercholesterolemia, 939-946
- Apolipoprotein(a) [apo(a)], metabolism of apo B-100 and, within plasma lipoprotein (a), 361-369
- Apolipoprotein A-I (apo A-I), effects on UC efflux, 1087-1094
- Apolipoprotein A-V (apo A-V)* gene, DNA variant sites in, association with CHD in Chinese, 568-572
- Apolipoprotein B (apo B)
- apo C-III bound to lipoproteins containing, and insulin resistance in Cherokee Indian youth, 180-187
 - production in laboratory opossums exhibiting high and low lipemic responses to dietary cholesterol and fat, 1075-1081
- Apolipoprotein B-48 (apo B-48)
- Lp particles containing, intestinal assembly and secretion in fasting state, 689-697
 - postprandial clearance, effect of LPL S447X variant, 1499-1503
- Apolipoprotein B-100 (apo B-100), metabolism of apo(a) and, within plasma lipoprotein (a), 361-369
- Apolipoprotein C-III (apo C-III)
- bound to apo B-containing lipoproteins, and insulin resistance in Cherokee Indian youth, 180-187
 - plasma, *see* Plasma apolipoprotein C-III
- Apolipoprotein E (apo E)
- influence of genotype on reliability of Friedewald formula in estimation of LDL-C concentrations, 1014-1019
 - recycling, physiological relevance, studies in primary mouse hepatocytes, 1309-1315
- AQP, *see* Aquaglyceroporins
- Aquaglyceroporins (AQP), and GlyK, regulations of, effects of TZDs in OLETF rats, 1282-1289
- Arg211His polymorphism of *GPR40* gene, contribution to variation of insulin secretory capacity in Japanese men, 296-299
- Aromatic ring-containing drugs, effect on carnitine biosynthesis with special regard to *p*-aminomethylbenzoic acid, 1582-1586
- Arrhythmia, outcomes in animal models, systematic review and meta-analysis of impact of ω -3 fatty acids, 1557-1565
- Arterial smooth muscle cells, human, effects of insulin, glucose, IGF-1, or growth hormone on formation and function of HA receptor CD44 on, 287-295
- Arterial stiffness and thickness, association with high serum pentosidine concentrations, in NIDDM, 345-350
- Arteriosclerosis obliterans, and IHD, association with hypoadiponectinemia, 653-656
- Artery compliance, effect of HF meal in subgroup of young healthy subjects, 1337-1344
- ASO, *see* Antisense oligonucleotides
- Aspartate aminotransferase (AST), *see* Plasma aspartate aminotransferase
- AST, *see* Plasma aspartate aminotransferase
- Asthma, PM women treated with inhaled corticosteroids for, effects of alendronate on BMD and bone metabolic markers in, 85-90
- Asymmetric dimethylarginine (ADMA), relationship to *S*-adenosylmethionine, *S*-adenosylhomocysteine, and endothelial function, in healthy human subject during experimental hyper- and hypohomocysteinemia, 351-360
- AT, *see* Adipose tissue
- Atherogenesis, endothelial dysfunction, and plasma adiponectin levels, effect of PKU diet, 1350-1355
- Atherogenicity, of LDL, effect of weight loss associated with reduced intake of CHO in premenopausal women, 1133-1141
- Atherogenic lipoprotein subfraction profile, in preeclamptic women with and without high TGs, 1504-1509
- Atherosclerosis
- apparent and non-apparent, in NIDDM, LDL size and subclasses as markers of, 227-234
 - novel risk factors, effect of rosiglitazone in patients with NIDDM and hypertension, 1236-1242
 - risk factors, levels of sICAM-1, sVCAM-1 and selectins in children and adolescents with, 1020-1026
 - subclinical, association with circulating sCD44, 139-141
- Athletes, middle-aged and young, versus sedentary men, effects of exercise at moderate and hard intensity on substrate oxidation, 1411-1419
- Atorvastatin
- low-dose, effect on circulating MCP-1 in patients with NIDDM complicated by hyperlipidemia, 1225-1229
 - and pravastatin and fenofibrate, comparative study of beneficial effects in types IIa and IIb hyperlipidemia patients, 677-681
 - in targeting CV risk in nondiabetic patients with MS, 1065-1074
- AT1R, *see* Angiotensin II AT1 receptor
- AT2R, *see* Angiotensin II AT2 receptor
- Atrasentan
- effect on hepatic insulin sensitivity associated with insulin signaling in Zucker fatty rats, 1515-1523
 - effects on metabolic responses in model of IR, 735-740
- Atrial natriuretic peptide (ANP), lipolytic pathways dependent on adrenergic receptor and, differential regulation in human SAT, 122-131
- Atropine, and electrical stimulation of vagal nerve, on insulin sensitivity, 579-583

- Autoantibodies, antipericyte, association with albuminuria and hypertension in NIDDM, 188-193
- Autogel, lanreotide, pharmacokinetics and population pharmacodynamic analysis, 1276-1281
- Autonomic function, cardiac, influence of nutrients in nondiabetic overweight subjects, 1290-1296
- Autonomic nervous system (ANS), cardiac, impaired response to glucose load in severe obesity, 966-974
- AvaII polymorphism, of *LDL-receptor* gene, and baseline LDL-C and Lp(a), influence on LDL-C response to pravastatin, 741-747
- Bacterial count, of aerobes and anaerobes, and serum lipids, effect of antibiotics on, 103-112
- BAECs, *see* Bovine aortic endothelial cells
- Baltimore Longitudinal Study of Aging, contribution of impaired glucose tolerance in subjects with metabolic syndrome, 542-547
- Basedow disease, effect of thyroid hormone on serum adiponectin levels in, 1461-1466
- BAT, *see* Brown adipose tissue
- BD, *see* Bipolar depression
- Beef intake, versus soy-based food supplements, effects on serum Lp-lipid profiles, CHO intake and nutrient intake ratios in older men who perform RT, 769-774
- Beta₂ (β_2) adrenergic receptor haplotype, and sex, effect on resting and terbutaline-stimulated whole body lipolysis, 492-499
- Beta (β) cell(s) (islets of Langerhans)
contribution of *GPR40* gene Arg211His polymorphism on functional variations of, in Japanese men, 296-299
function, effects of metformin on, 235-239
functional nicotinic receptors in, 247-254
gene expression in, and hyperglycemia prevention in Zucker diabetic fatty rats, effects of exercise training on, 1571-1581
- Beta (β)-thalassemia major, versus CDGP, hCG therapy in adolescent boys with, 15-23
- Bile duct ligation, and dimethylnitrosamine-induced liver cirrhosis, TNF- α -associated leptin expression in, 445-452
- Biliary cholesterol secretion, in patients with cerebrotendinous xanthomatosis, 335-344
- Biokinetics, of buccal spray insulin in IDDM patients, 930-934
- Bipolar depression (BD)
genetics and genomics, S1:11
linkage analysis, S1:11-12
overlap with MDD, S1:11
- Birth weight, and waist circumference, and serum hepatic enzyme levels, of 18-year old sons, associations with their father's and mother's BMI, 466-470
- Blood coagulation, exercise-induced activation of, effects of α -lipoic acid supplementation, 815-820
- Blood glucose, effect of vagal nerve electrical stimulation on, 581
- Blood glucose-lowering therapies, withdrawal of, and VLCD, effect on endogenous glucose production in obese NIDDM, 705-712
- Blood lactate, relationship to substrate utilization during exercise in NIDDM postmenopausal women, 1102-1107
- Blood lipids
effect of pioglitazone on, 24-32
endothelial activation and inflammation in prepubertal obese Turkish children, 1384-1389
and insulin sensitivity in overweight and obese girls, effects of aerobic exercise training, 1472-1479
see also Dyslipidemia; Hypercholesterolemia; Hyperlipidemia; Hypoalphalipoproteinemia; Lipemic responses; Postprandial lipemia (lipidemia)
- Blood lipoprotein (Lp), and insulin sensitivity in overweight and obese girls, effects of aerobic exercise training, 1472-1479
- Blood pressure (BP)
changes, and plasma catecholamine levels, during prolonged hyperinsulinemia, 391-396
and ECG data, effects of chronic GH and IGF-1 excess in young adults with pituitary gigantism, 1177
effects of soy-rich diet and stevioside in NIDDM rats, 1186
long-term and immediate effects of intranasal administration of insulin, 1356-1361
regulation in fructose-induced insulin resistance and hypertension, functional interaction of AT1R and AT2R in, 157-164
relation to high-sensitive CRP, TNF- α , lipids and insulin resistance before and after weight loss in obese children, 1155-1161
see also Diastolic blood pressure; Systolic blood pressure
- Blood viscosity, at multiple shear rates, influence of Lp, 764-768
- BMD, *see* Bone mineral density
- BMI, *see* Body mass and body mass index
- Body composition
and EE, effect of pioglitazone on, 24-32
and growth, pubertal alterations in spontaneous secretion and metabolic clearance of GH in overweight youth, 1374-1383
and metabolism, in male offspring of high-fat fed rats, 500-507
role in development of IR in primary renal disease, 821-828
- Body fat, *see* Adipose tissue
- Body mass and body mass index (BMI)
correlation to adiponectin-leptin ratio, in NIDDM, 281-286

- Body mass and body mass index (BMI) (continued)
correlation to plasma leptin levels, in children with short stature administered with insulin, 862-865
of middle-aged mothers and fathers, associations with 18-year-old son's waist circumferences, birth weight, and serum hepatic enzyme levels, 466-470
relationship to insulin resistance, soluble E-selectin, and serum lipid profile, in nonobese Japanese with NIDDM, 376-380
and sTNF-R, leptin, adiponectin and serum lipid profile in nonobese Japanese NIDDM patients, 1297-1301
and waist circumferences, elderly versus young men with similar and smaller, oxidative and nonoxidative glucose disposal in, 748-755
and WC, relationship between IMGU and excess adiposity, 1323-1328
- Body temperature, effects on tissue protein synthesis in rats, 841-847
- Body weight (BW)
in chronic blockade of NO synthesis, 483-484
control, and insulin sensitivity, effect of RVF containing high cyclo (his-pro), 1480-1489
effect of pioglitazone and insulinemia or sulfonylurea usage on, 28-29
effects of HX531 in KK-A^y mice, 574-576
effects of soy-rich diet and stevioside in NIDDM rats, 1186
effects of VLCFAs (policosanols) and phytosterols on, 511
and food consumption and adipose depots, effects of camostat, 621-622
and insulin sensitivity in overweight and obese girls, effects of aerobic exercise training, 1472-1479
of male offspring of high-fat fed rats, 503
see also Non-obese (lean) subjects; Obesity
- Bone loss, in castrated male mice, effect of *Puerariae radix*, 1536-1541
- Bone metabolic markers, in PM asthmatic women treated with inhaled corticosteroids, effects of alendronate versus alfacalcidol on, 85-90
- Bone mineral density (BMD), in PM asthmatic women treated with inhaled corticosteroids, effects of alendronate versus alfacalcidol on, 85-90
- Bone resorption, effect of simultaneous dietary administration of xylitol and ethanol, 548-551
- Bovine aortic endothelial cells (BAECs), effect of MET on intracellular production of ROS in, 829-834
- Boys
adolescent, with CDGP versus those with β -thalassemia major, hCG therapy in, 15-23
prepubertal, racial difference in circulating SHBG levels in, 91-96
with recent onset of obesity, effect of high-carbohydrate meal on postprandial thermogenesis and sympathetic nervous system activity in, 430-438
- BP, *see* Blood pressure
- Brain glucose, concentrations in poorly controlled DM, measurement by high-field magnetic resonance spectroscopy, 1008-1013
- Brain remodeling
adaptive structural plasticity in response to stress, S1:20-21
stress-induced changes in prefrontal cortex and amygdala, S1:21-22
- Brain stem, role in neurophysiologic modulation of obesity, 1209
- Brown adipose tissue (BAT)
accumulation, and UCP3 expression in skeletal muscle, effect of NO-1886 on HF diet-induced obesity, 1587-1592
role in neurophysiologic modulation of obesity, 1204
- Bu, *see* Gamma (γ)-butyrobetaine
- Buccal spray insulin, biokinetics of, in IDDM patients, 930-934
- BW, *see* Body weight
- C, *see* Cholesterol
- CAD, *see* Coronary artery disease
- Calcium channel blockers (CCBs), in targeting CV risk in nondiabetic patients with MS, 1065-1074
- Calcium-induced workload, and fatty acid supply, influence on myocardial substrate selection, 410-420
- Caloric restriction, effects on diurnal pattern of IGFBP-1 levels, in NIDDM, 275-280
- Camostat
effects on body weight, food consumption and adipose depots, 621-622
effects on dyslipidemia, insulin secretory defects and histological abnormalities of pancreas in genetically obese and diabetic rats, 619-627
effects on fasting serum TGs, cholesterol and FFAs, 623
effects on serum insulin, glucose response to IVGTT, IR and pancreatic insulin content, 623
- Captopril, effect on reflex increases in adrenal or lumbar SNA induced by hypoglycemia, 1679-1686
- Carbohydrate(s) (CHO)
intake of, and serum Lp-lipid profile and nutrient intake ratios, effects of beef versus soy-based food supplements in older men who perform RT, 769-774
labeled, administered at rest, role of labeled CO₂ production and oxidative versus nonoxidative disposal of, 1428-1434
reduced intake, associated with weight loss, effect on atherogenicity of LDL in premenopausal women, 1133-1141
supplementation, versus isoenergetic protein ingestion, combined with resistance training, effect on muscle fiber size and muscle strength, 151-156
see also Oxidation, CHO and specific carbohydrates

- Carbon dioxide (CO₂), production of labeled, and oxidative versus nonoxidative disposal of labeled CHO administered at rest, 1428-1434
- CARDIA, *see* Coronary Artery Risk Development in Young Adults
- Cardiac autonomic function, influence of nutrients in nondiabetic overweight subjects, 1290-1296
- Cardiac autonomic nervous system (ANS), impaired response to glucose load in severe obesity, 966-974
- Cardiac myocytes, cultured, effect of elevated glucose on protein synthesis in, 1453-1460
- Cardiorespiratory fitness, relation to normal WBC count in apparently healthy Korean men, 1448-1452
- Cardiovascular autonomic dysfunction, and insulin resistance, association with highly-sensitivity C-reactive protein, in NIDDM, 552-558
- Cardiovascular (CV) disease
- association with high serum pentosidine concentrations, in NIDDM, 345-350
 - effects on urinary excretion of glycine betaine and *N,N*-dimethylglycine, in patients attending a lipid clinic, 5-6
 - and NIDDM, development from obesity and MS, sex difference in subjects with normal glucose metabolism, 800-808
 - reciprocal relation to depression, S1:45-48
 - risk, *see* Risk, for CV disease
 - risk factors in healthy Korean women, association with circulating OPG levels, 49-54
- Carnitine
- biosynthesis, effect of aromatic ring-containing drugs with special regard to *p*-aminomethylbenzoic acid, 1582-1586
 - in fasting plasma and urine samples of patients attending a lipid clinic, 1-14
 - supplementation, effect on lipoprotein metabolism and LDL atherogenicity, 1135
- Carotid artery intimal-media thickness (IMT)
- in assessment of LDL size and subclasses as markers of atherosclerosis in NIDDM, 227-234
 - and lipid profile, in adults with growth hormone deficiency after long-term growth hormone replacement, 321-329
- Carotid intimal hyperplasia, and serum Hcy and smooth muscle proliferation, effect of rosiglitazone in high methionine-fed Sprague-Dawley rats, 645-652
- Catecholamine, *see* Plasma catecholamine
- CBG, *see* Corticosteroid-binding globulin
- CβS, *see* Cystathionine-β-synthase
- CCBs, *see* Calcium channel blockers
- CD44, formation and function in human arterial smooth muscle cells, effects of insulin, glucose, IGF-1, or growth hormone, 287-295
- CDGP, *see* Constitutional delayed growth and puberty
- Cellular insulin resistance, in Epstein-Barr virus-transformed lymphoblasts from young insulin-resistant Japanese men, 370-375
- Cerebral malaria, glucose production and plasma adiponectin in, 60-66
- Cerebrotendinous xanthomatosis (CTX), isotopomer spectral analysis of cholesterol synthesis intermediates in patients with, 335-344
- CHD, *see* Coronary heart disease
- Chemiluminescence assay, in vitro neutrophil oxidative burst, effects of propylthiouracil and thiamazole, 300-305
- Chennai Urban Rural Epidemiology Study, association of low levels of adiponectin with metabolic syndrome, 476-481
- Cherokee Indians, *see* Race and ethnicity
- CHH, *see* Chronic hypobaric hypoxia
- Children
- and adolescents, *see* Children, adolescents and
 - Cherokee Indians, apo C-III bound to apo B-containing lipoproteins and insulin resistance in, 180-187
 - with idiopathic short stature and growth hormone deficiency, leptin responses to insulin administration in, 862-865
 - obese, *see* Children, obese
- Children, adolescents and
- with obesity, hypertension and IDDM, levels of sICAM-1, sVCAM-1 and selectins, 1020-1026
 - relationship between fasting plasma ghrelin and metabolic parameters in, 925-929
- Children, obese
- high-sensitive CRP, TNF-α and cardiovascular risk factors before and after weight loss in, 1155-1161
 - insulin sensitivity indices of glucose and free fatty acid metabolism, in relation to serum lipids in, 397-402
 - and overweight girls, effects of aerobic exercise training on insulin sensitivity, body weight, body fat, adiponectin and inflammatory markers, 1472-1479
 - prepubertal Turkish, endothelial activation and inflammation in, 1384-1389
- Chinese subjects
- association between DNA variant sites in *apo A5* gene and CHD in, 568-572
 - association between lower serum free thyroxine levels and MS in, 1524-1528
- Chlamydia pneumoniae* infection, prevalence of, in relation to CRP levels, in subjects with hypoalphalipoproteinemia, 33-37
- CHO, *see* Carbohydrate
- Cholecystokinin, role in neurophysiologic modulation of obesity, 1210
- Cholesterol (C)
- accumulation, and macrophage scavenger receptor I expression, effect of iron loading in THP-1 mononuclear phagocytes, 453-459

- Cholesterol (C) (continued)**
and fasting serum TGs and FFAs, effects of camostat, 623
plasma, *see* Plasma cholesterol
unesterified, *see* Unesterified cholesterol
see also High-density lipoprotein-cholesterol;
Hypercholesterolemia; Low-density lipoprotein-cholesterol; Very-low-density lipoprotein-cholesterol
- Cholesterol absorption inhibitor, SCH 58235, and high-plant-sterol diet, effects on plant sterol absorption and plasma concentrations in hypercholesterolemic rats, 38-48**
- Cholesterol biosynthesis, and plasma lipid levels, effects of very long chain fatty acids (policosanols) and phytosterols, 508-814**
- Cholesterol efflux**
decreased, HDL-mediated, familial massive tendon xanthomatosis with, 1095-1101
endothelial, promoted by HDL APF, 1087-1094
in the presence of PCs in free form, 1091
- Cholesterol synthesis, intermediates of, isotopomer spectral analysis in cerebrotendinous xanthomatosis, 335-344**
- Chromium picolinate, effects of chronic treatment in uninephrectomized rats, 1243-1249**
- Chromosome 3p21, absence of fatty liver in FHBL linked to, 682-688**
- Chronic chromium picolinate treatment, effects in uninephrectomized rats, 1243-1249**
- Chronic hypobaric hypoxia (CHH), anorexia induced by, and RV hypertrophy, role in lactate transport and monocarboxylate transporter expression in rat muscle, 634-644**
- Chronic infections, prevalence of, in relation to CRP levels, in subjects with hypoalphalipoproteinemia, 33-37**
- Ciprofloxacin, effect on serum lipids and colonic microflora, 103-112**
- Circulating adiponectin, relation to cortisol, effect of CBG in men and women, 584-589**
- Circulating antipercyte autoantibodies (APAAAs), association with albuminuria and hypertension in NIDDM, 188-193**
- Circulating free fatty acids (FFAs), elevated concentrations of, effect on human SM gene expression, 952-959**
- Circulating insulin-like growth factor-1 (IGF-1), and IGF-BPs, genetic regulation of variation in human pedigrees, 975-981**
- Circulating insulin-like growth factor-binding protein (IGF-BPs), and IGF-1, genetic regulation of variation in human pedigrees, 975-981**
- Circulating insulin-like growth factor-binding protein 1 (IGFBP-1), altered response to nutritional deprivation in NIDDM, 275-280**
- Circulating monocyte chemoattractant protein-1 (MCP-1), effect of low-dose atorvastatin in patients with NIDDM complicated by hyperlipidemia, 1225-1229**
- Circulating osteoprotegerin (OPG), association with age, WHR, serum TC, and LDL-C levels in healthy Korean women, 49-54**
- Circulating soluble CD44 (sCD44), associations with sex, cardiovascular risk factors, and subclinical atherosclerosis, 139-141**
- Clinical course, variant, of 2 patients with neonatal intrahepatic cholestasis who have novel mutation of SLC25A13, 1615-1619**
- CMV, *see* Cytomegalovirus**
- CO₂, *see* Carbon dioxide**
- Coadministration, of raloxifene and low-dose simvastatin, effects on serum lipids in PM women with primary hypercholesterolemia, 939-946**
- Cobalamin, dose regimen for maximum Hey reduction in ESRD, 835-840**
- Cocaine transcript, role in neurophysiologic modulation of obesity, 1208-1209**
- Colonic microflora, and serum lipids, effect of antibiotics on, 103-112**
- Constitutional delayed growth and puberty (CDGP), versus β -thalassemia major, hCG therapy in adolescent boys with, 15-23**
- Coronary angioplasty, NIDDM subjects with CAD after, effect of rosiglitazone on clinical inflammatory responses in, 590-597**
- Coronary artery disease (CAD)**
association with fasting apo B-48 level in population without frank fasting hypertriglyceridemia, 1442-1447
in NIDDM, after coronary angioplasty, effect of rosiglitazone on clinical inflammatory responses in, 590-597
risk, *see* Risk, for CAD
risk factors for, and fibrinogen plasma concentration, 165-170
see also Atherosclerosis
- Coronary Artery Risk Development in Young Adults (CARDIA), 910-917**
- Coronary heart disease (CHD)**
association with SNPs of apo A5 gene, in Chinese, 568-572
LDL size and subclasses as markers of, in NIDDM, 227-234
- Corticosteroid-binding globulin (CBG), effect on circulating adiponectin-cortisol relationship, in men and women, 584-589**
- Corticosteroids, inhaled, effects of alendronate versus alfacalcidol on BMD and bone metabolic markers in PM asthmatic women treated with, 85-90**
- Corticosterone, *see* Plasma corticosterone**
- Corticotropin, *see* Plasma corticotropin**

- Corticotropin-releasing hormone (CRH), and glucocorticoid, combined deficiency, effect on counterregulatory responses after recurrent hypoglycemia in mice, 1259-1265
- Cortisol (hydrocortisone)
hyperleptinemia induced by, energy consumption in, 876-880
plasma, *see* Plasma cortisol
relation to circulating adiponectin, effect of CBG, in men and women, 584-589
serum, *see* Serum cortisol
- Cortisol, supplementation, effect on serum cortisol responses to physical stress, 657-668
- Counterregulatory deficits, after recurrent hypoglycemia, effect of combined CRH and glucocorticoid deficiency, 1259-1265
- CP, *see* C-peptides
- C-peptide (connecting peptides; CP), levels, and serum IL-6, association with plasma Hcy in NIDDM patients, 809-814
- Cranberry juice supplementation, effect on plasma antioxidant capacity and OxLDL levels in men, 856-861
- CrCl, *see* Creatinine clearance
- C-reactive protein (CRP)
or adiponectin, association with exercise training, 533-541
endothelial inflammation in prepubertal obese Turkish children, 1384-1389
high-sensitivity, *see* High-sensitivity C-reactive protein
and insulin resistance, reciprocal association with adipocytokines, in clinically healthy men, 439-444
and insulin sensitivity in overweight and obese girls, effects of aerobic exercise training, 1472-1479
levels, in relation to prevalence of chronic infections in hypoalphalipoproteinemia, 33-37
and Lp(a) and Hcy, levels in PM women, effects of types of HRT in PM women, 72-78
in patients with NIDDM and hypertension, effect of rosiglitazone, 1236-1242
plasma, *see* Plasma C-reactive protein
- Creatinine clearance (CrCl), role in development of IR in primary renal disease, 821-828
- CRH, *see* Corticotropin-releasing hormone
- CRP, *see* C-reactive protein
- CS, *see* Cytosolic sialidase
- C161→T polymorphism, in PPAR γ , association with insulin resistance in Hispanic and non-Hispanic white women, 1552-1556
- CTX, *see* Cerebrotendinous xanthomatosis
- CV disease, *see* Cardiovascular disease
- Cyanocobalamin, versus hydroxycobalamin, effects on plasma Hcy concentrations in ESRD, 1362-1367
- Cyclosporine, influence on LDL uptake in human lymphocytes, 1620-1625
- CYP2E1, *see* Cytochrome P4502E1
- Cystathionine- β -synthase (C β S), activity of, effect of rosiglitazone in high methionine-fed Sprague-Dawley rats, 649
- Cytochrome P4502E1 (CYP2E1) cells, mechanism of action of ethanol in down-regulation of 2,6-ST mRNA in, 729-734
- Cytokines
and glial function, relation to depression, S1:33-38
inflammatory, and adipocytokine levels, effects of diet and/or exercise in postmenopausal NIDDM women, 866-875
- Cytomegalovirus (CMV) infection, prevalence of, in relation to CRP levels, in subjects with hypoalphalipoproteinemia, 33-37
- Cytosolic sialidase (CS), activities and RSRs in rat liver, effects of chronic ethanol feeding, 1056-1064
- DAG, *see* Diacylglycerol
- Dehydroepiandrosterone sulfate (DHEA-S), *see* Plasma dehydroepiandrosterone sulfate; Serum dehydroepiandrosterone sulfate
- Depressive disorders
epidemiology, S1:6
genetics and genomics, S1:10-15
history, S1:5-6
humoral markers, S1:42-43
introduction, S1:2-4
and menopause, S1:51
morbid conditions associated with, S1:45-46
morphological changes in brain associated with, S1:42
neurobiological basis, S1:24-27
and neurosurgery, S1:28-32
pathophysiology, S1:6-7
pharmacotherapy, S1:7-8
postpartum, S1:50-51
during pregnancy, S1:50
reciprocal relation to CV disease, S1:45-48
relation to cytokines and glial function, S1:33-38
relation to genes and stress, S1:16-19
relation to glucocorticoids and mood disorders, S1:20-23
somatic markers, S1:41-42
subsyndromal depression, underdiagnosis and undertreatment in elderly, S1:39-44
symptoms, "sickness behavior" and cytokines, S1:34
in women, S1:49-52
see also specific disorders
- DGAT, *see* Diacylglycerol acyltransferase
- DHA, *see* Docosahexaenoic acid
- DHEA-S, *see* Dehydroepiandrosterone sulfate
- Diabetes mellitus (DM)
alloxan-induced, effects of FK-614 on peripheral glucose utilization and hepatic insulin extraction in dogs, 1250-1258
hepatic glucose production and hyperglycemia in, effects of ASO targeted against GCCR, 848-855

- Diabetes mellitus (DM) (continued)
poorly controlled, measurement of brain glucose concentrations by high-field magnetic resonance spectroscopy, 1008-1013
see also Gestational diabetes mellitus; Insulin-dependent diabetes mellitus; Maturity-onset diabetes of the young; Non-insulin-dependent diabetes mellitus; Streptozotocin-induced diabetes mellitus
- Diabetic microangiopathy, *see* Non-insulin-dependent diabetes mellitus microangiopathy
- Diabetic nephropathy, *see* Non-insulin-dependent diabetes mellitus nephropathy
- Diabetic retinopathy, *see* Non-insulin-dependent diabetes mellitus retinopathy
- Diabetogenic effects, of prolonged FFA exposure, susceptibility of young, LBW men to, 1398-1406
- Diacylglycerol (DAG), dietary, effects on serum TAGs in patient homozygous for complete lipoprotein lipase deletion, 67-71
- Diacylglycerol acyltransferase (DGAT), increased activity, association with triglyceride accumulation in tissues of diet-induced insulin-resistant hyperlipidemia, 403-409
- Diastolic blood pressure (BP), endothelial activation and inflammation in prepubertal obese Turkish children, 1384-1389
- Diet
benefits of metformin and, in women with PCOS and IR, 113-121
enriched with *cis*- or *trans*-FAs, comparison of immediate response to glucose and lipids in overweight individuals with differing FABP2 genotypes, 1652-1658
and/or exercise, effects on adipocytokine and inflammatory cytokine levels in postmenopausal NIDDM women, 866-875
high plant sterol-enriched, and SCH 58235, effects on plant sterol absorption and plasma concentrations in hypercholesterolemic rats, 38-48
PKU, effect on plasma adiponectin concentrations, endothelial dysfunction and atherogenesis, 1350-1355
see also Fasting; Food intake and entries beginning with terms: Dietary, Postprandial
- Dietary diacylglycerol (DAG), effects on serum TAGs in patient homozygous for complete lipoprotein lipase deletion, 67-71
- Dietary ethanol, and dietary xylitol, effect of simultaneous administration on bone resorption, 548-551
- Dietary modification, effect on lipoprotein metabolism and LDL atherogenicity, 1135
- Dietary obesity, phenotype-based treatment of, differential effects of fenofibrate, 421-429
- Dietary protein, source of, effects on serum Lp-lipid profiles, CHO intake and nutrient intake ratios, in older men who perform RT, 769-774
- Dietary *trans*-fatty acids (FA), effects on adipocyte plasma membrane fatty acid composition and insulin sensitivity, 240-246
- Dietary xylitol, and ethanol, effect of simultaneous administration on bone resorption, 548-551
- Diet-induced insulin-resistant hyperlipidemia, triglyceride accumulation in tissues of, association with increased diacylglycerol acyltransferase activity, 403-409
- D-dimer, role in assessment of fibrinogen metabolism in the lungs, 271-274
- Dimethylarginine, *see* Asymmetric dimethylarginine
- N,N*-Dimethylglycine, and Hcy, and glycine betaine, in patients attending a lipid clinic, 1-14
- Dimethylguanidine, *see* Metformin
- Dimethylnitrosamine (DMN), and bile duct ligation-induced liver cirrhosis, TNF- α -associated leptin expression in, 445-452
- Disabilities, associated with aging process, S1:40
- Disability improvement, after stroke, effect of PC and NTIS, 702
- DM, *see* Diabetes mellitus
- DMN, *see* Dimethylnitrosamine
- DNA mutations, in apo A5 gene, association with CHD in Chinese, 568-572
- Docosahexaenoic acid (DHA), effects on vascular reactivity in healthy adult rats, 1600-1609
- Doppler echocardiography, cardiac effects of chronic GH and IGF-1 excess in young adults with pituitary gigantism, 1177-1178
- Dose, of cobalamin for maximum Hcy reduction in ESRD, 835-840
- Dose-response effect, of differing volumes of resistance exercise bouts on postprandial lipemia, 756-763
- Dyslipidemia
and insulin secretory defects and histological abnormalities of pancreas, effects of synthetic PI camostat in genetically obese and diabetic rats, 619-627
see also Hypercholesterolemia; Hyperlipidemia; Hypoalphalipoproteinemia; Postprandial lipemia (lipidemia)
- Dysphagia, effect on nutrition status, 701
- EBV, *see* Epstein-Barr virus
- ecNOS, *see* Endothelial constitutive nitric oxide synthase
- ECs, *see* Endothelial cells
- EE, *see* Energy expenditure
- EGP, *see* Endogenous glucose production
- Eicosapentaenoic acid (EPA), effects on vascular reactivity in healthy adult rats, 1600-1609
- Electrocardiography, and BP, effects of chronic GH and IGF-1 excess in young adults with pituitary gigantism, 1177

- Endogenous glucose production (EGP)
effect of 2-day VLED on SM insulin sensitivity in obese NIDDM patients on insulin therapy, 1669-1678
effect of VLCD and withdrawal of blood glucose-lowering therapies in obese NIDDM, 705-712
postprandial control of, effects of repaglinide and gliclazide on, 79-84
- Endothelial activation, and inflammation, in prepubertal obese Turkish children, 1384-1389
- Endothelial cells (ECs), aortic, effect of MET on intracellular production of ROS in, 829-834
- Endothelial cholesterol efflux, promoted by HDL APF, 1087-1094
- Endothelial constitutive nitric oxide synthase (ecNOS), genotype, possible interactions with alcohol drinking and walking time for high serum uric acid levels among Japanese, 1302-1308
- Endothelial dysfunction, atherogenesis, and plasma adiponectin levels, effect of PKU diet, 1350-1355
- Endothelial function
effects of red wine consumption on insulin resistance and, in NIDDM, 306-313
relationship to S-adenosylmethionine, S-adenosylhomocysteine, and asymmetric dimethylarginine, in healthy human subject during experimental hyper- and hypohomocysteinemia, 351-360
- Endothelial nitric oxide synthase (eNOS), content in AT from obese and lean African American and white American women, 1368-1373
- Endothelin (ET) antagonism
with atrasentan, effect on hepatic insulin sensitivity associated with insulin signaling in Zucker fatty rats, 1515-1523
effects on metabolic responses in model of IR, 735-740
- Endothelium, monocyte adhesion to, relation to hypercholesterolemia and MS in nondiabetic subjects, 1467-1471
- End-stage renal disease (ESRD)
altered relationship of body fat and plasma adiponectin in, 330-334
cobalamin dose regimen for maximum reduction of Hcy in, 835-840
effects of hydroxycobalamin versus cyanocobalamin on plasma Hcy in, 1362-1367
- Endurance running, effects on plasma OPG and (s)RANKL, 935-938
- Endurance training
effects on substrate oxidation in middle-aged and young athletes versus sedentary men, 1411-1419
insulin action in response to, association with PPAR γ 2 Pro12Ala genotype in men and women, 97-102
- Energy (caloric) restriction, effects on diurnal pattern of IGFBP-1 levels, in NIDDM, 275-280
- Energy expenditure (EE), and body composition, effect of pioglitazone on, 24-32
- eNOS, *see* Endothelial nitric oxide synthase
- EPA, *see* Eicosapentaenoic acid
- Epstein-Barr virus (EBV)-transformed lymphoblasts, cellular insulin resistance in, 370-375
- ERK 1/2, *see* Extracellular signal-regulated kinase 1/2
- Erythrocyte deformability, beneficial effects of pravastatin, atorvastatin and fenofibrate in types IIa and IIb hyperlipidemia patients, 677-681
- E-selectin, *see* Soluble E-selectin
- ESRD, *see* End-stage renal disease
- Estradiol, endogenous, relation to serum adiponectin levels, 1610-1614
- Estrogen, natural transdermal versus oral synthetic, effect on Lp(a), Hcy, and CRP levels in PM women, 72-78
- ET, *see* Endothelin
- Ethanol, *see* Alcohol
- Euglycemic-hyperinsulinemic clamping, effects of FFAs on plasma resistin and insulin resistance during, 1143-1144
- Exercise
aerobic, effects on insulin sensitivity, body weight, body fat, adiponectin and inflammatory markers in overweight and obese girls, 1472-1479
association with levels of C-reactive protein or adiponectin, 533-541
blood coagulation activation induced by, effects of α -lipoic acid supplementation, 815-820
and/or diet, effects on adipocytokine and inflammatory cytokine levels in postmenopausal NIDDM women, 866-875
effects on hyperglycemia prevention in Zucker diabetic fatty rats and gene expression in insulin-sensitive tissues determined by high-density oligonucleotide microarray analysis, 1571-1581
endurance, *see* Endurance training
glycogen repletion in SM after, effect of impaired glucose uptake in insulin-treated STZ-DM fasted rats, 1420-1427
at moderate and hard intensity, effects on substrate oxidation in middle-aged and young athletes versus sedentary men, 1411-1419
oxidation of combined glucose and sucrose ingestion during, 610-618
relationship between blood lactate concentration and substrate utilization during, in NIDDM postmenopausal women, 1102-1107
repeated sprint, influence of allopurinol on urinary purine loss after, 1269-1275
resistance, *see* Resistance training
substrate availability and transcriptional regulation of metabolic genes in human SM during recovery from, 1048-1055
- Exercise bouts, single versus multiple, effect on glucose control in women with NIDDM, 989-994
- Exercise training, *see* Exercise

- Experimental hyperhomocysteinemia, relationship between *S*-adenosylmethionine, *S*-adenosylhomocysteine and asymmetric dimethylarginine, in healthy human subject during, 351-360
- Experimental hypohomocysteinemia, relationship between *S*-adenosylmethionine, *S*-adenosylhomocysteine and asymmetric dimethylarginine, in healthy human subject during, 351-360
- Extracellular signal-regulated kinase 1/2 (ERK 1/2), activity, and protein synthesis in cultured cardiac myocytes, effect of elevated glucose, 1453-1460
- F, *see* Fructose
- FA(s), *see* Fatty acid(s)
- FABP2, *see* Fatty acid-binding protein 2
- Factor V *Leiden* gene mutation, and high factor VIII and high PAI activity, etiologies for sporadic miscarriage, 1345-1349
- Factor VIII (FVIII)
and factor V *Leiden* gene mutation and high PAI activity, etiologies for sporadic miscarriage, 1345-1349
serum, *see* Serum factor VIII
- Familial hypobetalipoproteinemia (FHBL), linked to chromosome 3p21, absence of fatty liver in, 682-688
- Familial massive tendon xanthomatosis, with decreased HDL-mediated cholesterol efflux, 1095-1101
- Farnesyl pyrophosphate (FPP), role in pitavastatin transactivation of human serum *PON1* gene, 142-150
- Fasting
effect of GH on glucose production and glycogenolysis, 604-609
intestinal assembly and secretion of apo B48-containing Lp particles in, 689-697
steatosis induced by, effect of transgenic expression of mutant PPAR γ in liver, 1490-1498
see also entries beginning with term: Fasting
- Fasting apolipoprotein B-48 (apo B-48), levels, association with CAD in population without frank fasting hypertriglyceridemia, 1442-1447
- Fasting blood glucose (FBG)
effect of HF meal in subgroup of young healthy subjects, 1340
effects of soy-rich diet and stevioside in NIDDM rats, 1186
- Fasting hypertriglyceridemia, absence of frank abnormality, association between fasting apo B-48 levels and CAD, 1442-1447
- Fasting plasma free fatty acid (FFA), effects of soy-rich diet and stevioside in NIDDM rats, 1186
- Fasting plasma ghrelin, relationship to metabolic parameters in children and adolescents, 925-929
- Fasting plasma glucose (FPG)
effect of 2-day VLED on SM insulin sensitivity in obese NIDDM patients on insulin therapy, 1669-1678
and insulin concentration after 2-day VLCD, 708
- Fasting plasma insulin, predicts hepatic fat content in healthy individuals with normal ALT concentrations, 1566-1570
- Fasting serum free fatty acids, association with insulin secretion, effect of westernized lifestyle in Japanese men, 713-718
- Fasting serum insulin, spectrum of insulin sensitivity in Korean population, 1644-1651
- Fasting serum triglyceride(s) (TGs), and cholesterol and FFAs, effects of camostat, 623
- FAT/CD36 expression, effect of 2-day VLED on SM insulin sensitivity in obese NIDDM patients on insulin therapy, 1669-1678
- Fatty acid(s) (FAs)
chain length, spatial configuration and degree of unsaturation, short-term effects on glucagon secretion, 1329-1336
changes in metabolism of, induced by high-fat ketogenic diet despite transient ketosis, 1127-1132
trans-, effects on adipocyte plasma membrane fatty acid composition and insulin sensitivity, 240-246
effects on vascular reactivity in healthy adult rats, 1600-1609
exogenous, effect on intestinal assembly and secretion of apo B48-containing Lp particles in fasting state, 689-697
cis- or *trans*-FAs, meals enriched with, comparison of immediate response to glucose and lipids in overweight individuals with differing FABP2 genotypes, 1652-1658
and lipids, hepatic, physiological relevance of apoE recycling in primary mouse hepatocytes, 1311-1312
supply of, and calcium-induced workload transitions, influence on myocardial substrate selection, 410-420
 ω -3, systematic review and meta-analysis of impact on selected arrhythmia outcomes in animal models, 1557-1565
uptake and oxidation in muscle of high fat-fed rats, effect of short-term leptin treatment, 1218-1224
- Fatty acid-binding protein 2 (FABP2), genotype, comparison of immediate response to meals enriched with *cis*- or *trans*-FAs in overweight individuals with differing, 1652-1658
- Fatty liver, absence in FHBL linked to chromosome 3p21, 682-688
- FBG, *see* Fasting blood glucose
- FDP, *see* Fibrin(ogen) degradation products
- Females, *see* Sex; Women
- Fenofibrate
differential effects in phenotype-based treatment of dietary obesity, 421-429
pravastatin and atorvastatin, comparative study of beneficial effects in types IIa and IIb hyperlipidemia patients, 677-681
in targeting CV risk in nondiabetic patients with MS, 1065-1074

- Ferritin, *see* Serum ferritin
- FFAs, *see* Nonesterified fatty acids (free fatty acids; NEFAs)
- FHBL, *see* Familial hypobetalipoproteinemia
- Fibrinogen
- levels, beneficial effects of pravastatin, atorvastatin and fenofibrate in types IIa and IIb hyperlipidemia patients, 677-681
 - metabolism in the lungs, 271-274
 - in patients with NIDDM and hypertension, effect of rosiglitazone, 1236-1242
 - plasma, *see* Plasma fibrinogen
- Fibrin(ogen) degradation products (FDP), in assessment of fibrinogen metabolism in lungs, 271-274
- Fibrinolysis, effects of α -lipoic acid, 817
- Fish oil, systematic review and meta-analysis of impact on selected arrhythmia outcomes in animal models, 1557-1565
- FK-614, effects on peripheral glucose utilization and hepatic insulin extraction in alloxan-induced diabetic dogs, 1250-1258
- Flavonoid-rich cranberry juice supplementation, effect on plasma antioxidant capacity and OxLDL levels in men, 856-861
- Fluvastatin, sensitivity of angiotensin II-induced oxidative burst to, in neutrophils of hypercholesterolemic patients, 1147-1154
- Food intake
- and body weight and adipose depots, effects of camostat, 621-622
 - effects of VLCFAs (policosanols) and phytosterols on, 511
 - in glucocorticoid-induced hyperleptinemia, 876-880
 - influence on cardiac autonomic function in nondiabetic overweight subjects, 1290-1296
 - restriction, effects on body weight and plasma leptin levels in KK-A^y mice, 576
 - see also* Diet
- FPG, *see* Fasting plasma glucose
- FPP, *see* Farnesyl pyrophosphate
- Framingham Heart Study, genome-wide search for genes affecting serum uric acid levels, 1435-1441
- Free fatty acids, *see* Nonesterified fatty acids (free fatty acids; NEFAs)
- Friedewald formula, reliability in estimation of LDL-C concentrations, influence of apo E genotype, 1014-1019
- Fructose (F), and sucrose, hepatic adaptations to, 1189-1201
- Fructose-induced insulin resistance, and hypertension, functional interaction of AT1R and AT2R in, 157-164
- Fuel oxidation
- during human walking, 793-799
 - whole body, of male offspring of high-fat fed rats, 500-507
- Functional nicotinic receptors, in pancreatic β cells, 247-254
- Functional polymorphism, of Z-DNA-forming motif of promoter of *SLC11A1* gene, association with IDDM in Japanese, 628-633
- Fungata Study, 669-676
- fVIII, *see* Factor VIII
- Gal β 1, 4GlcNAc α 2,6-sialyltransferase messenger RNA (2,6-ST mRNA), down-regulation in human liver cell lines, effect of ethanol exposure, 729-734
- Gamma (γ)-butyrobetaine (Bu), unlabeled, inhibitory effect of PAMBA on conversion of, 1584-1585
- Ganglioside sialidase, activities and RSRs in rat liver, effects of chronic ethanol feeding, 1056-1064
- GCCR, *see* Glucocorticoid receptor
- GCR, *see* Glucagon receptor gene
- Gene expression, *see* Messenger ribonucleic acid
- Genetic linkage
- genome-wide search for genes affecting serum uric acid, 1435-1441
 - of glucagon receptor gene with NIDDM in Italians, 786-787
- Genetic regulation, of variation of circulating IGF-1, IGF-BPs and leptin in human pedigrees, 975-981
- Genotype
- ACP₁, relation to serum glucose concentration in healthy adult subjects, 891-894
 - apo E, influence on reliability of Friedewald formula in estimation of LDL-C concentrations, 1014-1019
 - ecNOS, possible interactions with alcohol drinking and walking time for high serum uric acid levels among Japanese, 1302-1308
 - FABP2, comparison of immediate response to meals enriched with *cis*- or *trans*-FAs in overweight individuals with differing, 1652-1658
 - PPAR γ 2 Pro12Ala, association with endurance training-induced changes in insulin response to oral glucose, in men and women, 97-102
- Gestational diabetes mellitus (GDM)
- and NIDDM, *IPF-1/MODY4* gene missense mutation in Italian family with, 983-988
 - women with and without, MS at follow-up in index pregnancy, 1115-1121
- GH, *see* Growth hormone
- GHD, *see* Growth hormone deficiency
- Ghrelin
- fasting plasma, *see* Fasting plasma ghrelin
 - plasma, *see* Plasma ghrelin
 - role in neurophysiologic modulation of obesity, 1209-1210
- GIP, *see* Glucose-dependent insulinotropic polypeptide
- Glial function, and cytokines, relation to depression, S1:33-38
- Gliclazide, and repaglinide, effects on postprandial control of EGP, 79-84
- GLP-1, *see* Glucagon

- Glucagon (GLP-1)
 plasma, *see* Plasma glucagon
 role in neurophysiologic modulation of obesity, 1210-1211
- Glucagon receptor gene (GCR), genetic linkage with NIDDM in Italians, 786-787
- Glucagon response, impact on postprandial hyperglycemia in men with impaired glucose tolerance and NIDDM, 1168-1173
- Glucagon secretion, short-term effects of FAs as influenced by their chain length, spatial configuration and degree of unsaturation, 1329-1336
- Glucocorticoid receptor (GCCR), antisense oligonucleotide against, effects on hepatic glucose production and hyperglycemia in DM mice, 848-855
- Glucocorticoid(s)
 and aminergic receptors, neurobiological basis of depression, S1:26
 and CRH, combined deficiency, effect on counterregulatory responses after recurrent hypoglycemia in mice, 1259-1265
 hyperleptinemia induced by, energy consumption in, 876-880
 relation to depression and mood disorders, S1:20-23
- Gluconeogenesis
 and glycogenolysis, effect of GH in fasting healthy volunteers, 604-609
 and plasma adiponectin, in patients infected with *Plasmodium falciparum*, 60-66
- Glucophage, *see* Metformin
- Glucose
 brain, *see* Brain glucose
 combined ingestion with sucrose, oxidation during exercise, 610-618
 elevated levels, effect on protein synthesis in cultured cardiac myocytes, 1453-1460
 endothelial activation and inflammation in prepubertal obese Turkish children, 1384-1389
 and free fatty acid metabolism, insulin sensitivity indices of, in relation to serum lipids, in obese children and adolescents, 397-402
 and insulin, IGF-1, or growth hormone, effects on formation and function of HA receptor CD44, on human arterial smooth muscle cells, 287-295
 and lipid metabolism after liver transplantation in inbred rats, 881-890
 and lipid oxidation rates after 2-day VLCD, 709
 and lipids, comparison of immediate response to meals enriched with *cis*- or *trans*-FAs in overweight individuals with differing FABP2 genotype, 1652-1658
 load, cardiac ANS response to, in severe obesity, 966-974
 plasma, *see* Plasma glucose
 production of, *see* Gluconeogenesis
 response to IVGTT, and serum insulin, IR and pancreatic insulin content, effects of camostat, 623
- Glucose (continued)
 serum, *see* Serum glucose
 see also Gestational diabetes mellitus; Glucose disposal; Glucose homeostasis; Glucose metabolism; Glucose-6-phosphate dehydrogenase; Glucose tolerance; Glucose transport; Glucose transporter 4; Glucose turnover; Glucose uptake; Glucose usage; Glucose utilization; Insulin-dependent diabetes mellitus; Maturity-onset diabetes mellitus of the young; Non-insulin-dependent diabetes mellitus; Streptozotocin-induced diabetes mellitus and entries beginning with prefixes: Gluc-, Glyc-
- Glucose control, *see* Glycemic control
- Glucose-dependent insulintropic polypeptide (GIP), role in neurophysiologic modulation of obesity, 1210-1211
- Glucose disposal
 effect of 2-day VLED on SM insulin sensitivity in obese NIDDM patients on insulin therapy, 1669-1678
 oxidative and nonoxidative, in elderly versus younger men with similar and smaller BMI and waist circumferences, 748-755
- Glucose homeostasis, in fructose-induced insulin resistance and hypertension, functional interaction of AT1R and AT2R in, 157-164
- Glucose metabolism
 and body weight control, effect of HCS plus RVE, 1480-1489
 and insulin sensitivity
 in offspring rats, effects of high saturated versus high-CHO diet during pregnancy and lactation, 1319
 in transgenic rabbits, effect of high LPL activity, 132-138
 whole-body, *see* Whole-body glucose metabolism
- Glucose-6-phosphate dehydrogenase (G6PDase), supramolecular complex with TALase, retrograde trafficking in human neutrophils during pregnancy, 1027-1033
- Glucose production, *see* Endogenous glucose production; Gluconeogenesis
- Glucose tolerance
 effect of modest lifestyle intervention in obese African Americans, 960-965
 effects of metformin on, 235-239
 impaired, and NIDDM, impact of glucagon response on postprandial hyperglycemia in men with, 1168-1173
- Glucose transport
 and Akt phosphorylation, effect of continually high insulin levels in human myoblasts, 1687-1693
 basal and insulin-stimulated, in human omental versus SC adipocytes, 781-785
 and insulin receptor activation, insulin-stimulated, analysis in cultured skeletal muscle cells from obese subjects, 598-603
 in muscle, and whole-body insulin sensitivity, effect of selective angiotensin II receptor antagonism on hypertensive TG(mREN2)27 rats, 1659-1668

- Glucose transporter 4 (GLUT4)
in human omental versus SC adipocytes, 781-785
translocation of, effect of 2-day VLED on SM insulin sensitivity in obese NIDDM patients on insulin therapy, 1669-1678
- Glucose turnover, during euglycemic-hyperinsulinemic clamping, effects of FFAs on plasma resistin and insulin resistance, 1144
- Glucose uptake
effect of p38 MAP kinase inhibition in insulin-resistant 3T3-L1 adipocytes, 895-901
impaired, effect on postexercise glycogen repletion in SM of insulin-treated STZ-DM fasted rats, 1420-1427
and insulin sensitivity in offspring rats, effects of high saturated versus high-CHO diet during pregnancy and lactation, 1319
- Glucose usage, in SM before and after exercise in insulin-treated STZ-DM fasted rats, 1424-1425
- Glucose utilization, peripheral, and hepatic insulin extraction, effects of FK-614 in alloxan-induced diabetic dogs, 1250-1258
- α (alpha)-glucosidase inhibitors, voglibose and acarbose, effects on postprandial hyperglycemia and subjective abdominal symptoms, 387-390
- GLUT4, *see* Glucose transporter 4
- Glycated (glycosylated) hemoglobin, *see* Hemoglobin A_{1c}
- Glycemic control, effect of single versus multiple bouts of exercise in women with NIDDM, 989-994
- Glycerol
and nonesterified fatty acids, lactate, triglycerides and hormones after 2-day VLCD, 709
see also Diacylglycerol; Serum triacylglycerols
- Glycerolipid synthesis, and secretion, by primary hepatocytes in culture, relevance of apoE recycling, 1312
- Glycerol kinase (GlyK), and AQP, regulations of, effects of TZDs in OLETF rats, 1282-1289
- Glycerol Ra, after 2-day VLCD, 709
- Glycine betaine, and Hcy and *N,N*-dimethylglycine, in patients attending a lipid clinic, 1-14
- Glycogenolysis, and glucose production, effect of GH in fasting healthy volunteers, 604-609
- Glycogen repletion, postexercise, effect of impaired glucose uptake in SM of insulin-treated STZ-DM fasted rats, 1420-1427
- Glycogen synthase fractional velocity, in SM before and after exercise in insulin-treated STZ-DM fasted rats, 1425
- Glycogen synthesis, and content, in offspring rats, effects of high saturated versus high-CHO diet during pregnancy and lactation, 1319-1320
- GlyK, *see* Glycerol kinase
- G6PDase, *see* Glucose-6-phosphate dehydrogenase
- GPR40 gene, Arg211His polymorphism, contribution to variation of insulin secretory capacity in Japanese men, 296-299
- Growth hormone (GH)
altered spontaneous secretion and metabolic clearance in overweight prepubertal and pubertal youth, 1374-1383
effect on glucose production and glycogenolysis, in fasting healthy volunteers, 604-609
human, *see* Human growth hormone
and IGF-1, chronic excess, cardiac and metabolic effects in young adults with pituitary gigantism, 1174-1180
and/or sex steroids, effect on whole-body protein turnover in healthy aged men and women, 1162-1167
- Growth hormone deficiency (GHD)
carotid artery IMT and lipid profile in, after long-term growth hormone replacement, 321-329
leptin responses to insulin administration in children with, 862-865
- Growth hormone replacement, long-term, carotid artery IMT and lipid profile in growth hormone deficiency after, 321-329
- G82S and G1704T polymorphisms of *RAGE* gene, association with microalbuminuria, in Japanese NIDDM patients, 488-491
- G1704T and G82S polymorphisms of *RAGE* gene, association with microalbuminuria, in Japanese NIDDM patients, 488-491
- Gut peripheral signals, role in neurophysiologic modulation of obesity, 1209-1211
- HAART, *see* Highly active antiretroviral therapy
- HAD, *see* High alcohol dehydrogenase
- 3-HB, *see* 3-Hydroxy-butylate
- HbA_{1c}, *see* Hemoglobin A_{1c}
- HBP, *see* High blood pressure
- HC, *see* Hypercholesterolemia
- hCG, *see* Human chorionic gonadotropin
- HCHF diet, *see* High-cholesterol and high-fat diet
- HCS, *see* High cyclo (his-pro) content in soy protein hydrolysate
- Hcy, *see* Homocysteine
- HDL, *see* High-density lipoprotein(s)
- HDL-C, *see* High density lipoprotein-cholesterol
- Heart
myofibrillar and sarcoplasmic protein synthesis in, effect of long-term alcohol administration on, 212-219
see also entries beginning with element: Card-
Helicobacter pylori infection, prevalence of, in relation to CRP levels, in subjects with hypoalphalipoproteinemia, 33-37
- Hemodynamic parameters, in chronic blockade of NO synthesis, 483-484
- Hemodynamics, effect of HF meal in subgroup of young healthy subjects, 1340
- Hemoglobin A_{1c} (glycated hemoglobin; HbA_{1c})
association with circulating APAAs in NIDDM, 188-193
endothelial activation and inflammation in prepubertal obese Turkish children, 1384-1389

- Hemostasis, effects of α -lipoic acid, 817
- Hepatic denervation, effect on glucose and lipid metabolism after liver transplantation in inbred rats, 881-890
- Hepatic enzyme, serum levels, and waist circumference, and birth weight, of 18-year old sons, associations with their father's and mother's BMI, 466-470
- Hepatic fat content, prediction in healthy individuals with normal ALT concentrations, role of hyperinsulinemia, 1566-1570
- Hepatic glucose production, and hyperglycemia, effects of ASO targeted against GCCR in DM mice, 848-855
- Hepatic glucose uptake, unique regulation by fructose, 1191
- Hepatic insulin extraction
and peripheral glucose utilization, effects of FK-614 in alloxan-induced diabetic dogs, 1250-1258
and posthepatic insulin clearance rate, in proportion to insulin resistance, in HIV infection with lipodystrophy, 171-179
- Hepatic insulin sensitivity (S_i), *see* Insulin sensitivity, hepatic
- Hepatic lipid content (HLC), inverse correlation with serum adiponectin concentration, in Japanese NIDDM subjects, 775-780
- Hepatic lipids, role in liver adaptations to sucrose and fructose, 1195
- Hepatocytes, mouse, physiological relevance of apoE recycling, 1309-1315
- HepG2 cells, wild-type, mechanism of action of ethanol in down-regulation of 2,6-ST mRNA in, 729-734
- Herpes simplex virus 1 infection, prevalence of, in relation to CRP levels, in subjects with hypoalphalipoproteinemia, 33-37
- HF diet, *see* High-fat diet
- HGA, *see* Homogentisic acid
- hGH, *see* Human growth hormone
- High alcohol dehydrogenase (HAD) cells, mechanism of action of ethanol in down-regulation of 2,6-ST mRNA in, 729-734
- High blood pressure (HBP), association with cellular insulin resistance in EBV-transformed lymphoblasts from young Japanese men, 370-375
- High-carbohydrate (CHO) diet
effect on postprandial thermogenesis and sympathetic nervous system activity, in boys with recent onset of obesity, 430-438
and high MUFA diet, TG response to, correlation to variations in plasma apo C-III levels, 1390-1397
versus high saturated fat diet during pregnancy and lactation, effects on insulin sensitivity in offspring rats, 1316-1322
- High-cholesterol and high-fat (HCHF) diet, LDL apo B production in laboratory opossums exhibiting high and low lipemic responses to, 1075-1081
- High cyclo (his-pro) content in soy protein hydrolysate (HCS), plus RVF, effect on insulin sensitivity and body weight control, 1480-1489
- High-density lipoprotein(s) (α -lipoprotein; HDL)
APF, effects on UC efflux, 1087-1094
decreased cholesterol efflux mediated by, familial massive tendon xanthomatosis with, 1095-1101
- High-density lipoprotein-cholesterol (HDL-C;
 α -lipoprotein-cholesterol)
correlation with adiponectin-leptin ratio, in NIDDM, 281-286
low levels of, *see* Hypoalphalipoproteinemia
serum, *see* Serum high-density lipoprotein-cholesterol
- High-density oligonucleotide microarray analysis, gene expression in insulin-sensitive tissues as determined by, effects of exercise training on hyperglycemia in Zucker diabetic fatty rats, 1571-1581
- High-fat (HF) diet
effect of short-term leptin treatment on FA uptake and oxidation in muscle of rats fed with, 1218-1224
effects on postprandial TGs in primary renal disease, 821-828
effect on vascular compliance in subgroup of young healthy subjects, 1337-1344
maternal, effect on body composition and metabolism, in male offspring rats, 500-507
obesity induced by, effect of NO-1886 on expression of UCP3 in skeletal muscle and fat accumulation, 1587-1592
steatosis induced by, effect of transgenic expression of mutant PPAR γ in liver, 1490-1498
- High-fat ketogenic diet, induces marked changes in fatty acid metabolism despite transient ketosis, 1127-1132
- Highly active antiretroviral therapy (HAART), PI-based, effect on thigh SAT gene expression, 561-567
- High methionine diet, effect of rosiglitazone on serum Hcy, smooth muscle proliferation and intimal hyperplasia in Sprague-Dawley rats, 645-652
- High-monounsaturated fatty acid (MUFA) diet, and high-CHO diet, TG response to, correlation to variations in plasma apo C-III levels, 1390-1397
- High saturated fat diet, versus high CHO diet during pregnancy and lactation, effects on insulin sensitivity in offspring rats, 1316-1322
- High-sensitivity C-reactive protein (CRP)
association with insulin resistance and cardiovascular autonomic dysfunction, in NIDDM, 552-558
in patients with NIDDM complicated with hyperlipidemia, effect of low-dose atorvastatin, 1225-1229
and TNF- α and cardiovascular risk factors before and after weight loss in obese children, 1155-1161
- Hispanic subjects, and non-Hispanic white women, association between C161 \rightarrow T polymorphism in PPAR γ and insulin resistance, 1552-1556
- Histological abnormalities, of pancreas, and dyslipidemia and insulin secretory defects, effects of synthetic PI camostat in genetically obese and diabetic rats, 619-627
- HIV, *see* Human immunodeficiency virus

- HLC, *see* Hepatic lipid content
- HMG-CoA, *see* Hydroxymethyl glutaryl coenzyme A
- HNE, *see* α,β -unsaturated aldehyde, 4-hydroxy-2-nonenal
- HOMA, *see* Homeostasis model assessment
- HOMA_{IR}, *see* Homeostasis model assessment insulin resistance
- Homeostasis model assessment (HOMA), spectrum of insulin sensitivity in Korean population, 1644-1651
- Homeostasis model assessment (HOMA)- β , relation to CrCl in patients with primary renal disease, 821-828
- Homeostasis model assessment insulin resistance (HOMA_{IR}) index
- contribution of *GPR40* gene Arg211His polymorphism on variation of, in Japanese men, 296-299
 - effect of HF meal in subgroup of young healthy subjects, 1340
 - limitation of the validity of, 206-211
 - relation to parameters of body composition in patients with primary renal disease, 821-828
- Homocysteine (Hcy)
- effect of methimazole-induced hypothyroidism, 460-465
 - and glycine betaine and *N,N*-dimethylglycine, in patients attending a lipid clinic, 1-14
 - and Lp(a) and CRP, levels in PM women, effects of types of HRT, 72-78
 - maximum reduction in ESRD, dose regimen of cobalamin for, 835-840
 - in patients with NIDDM and hypertension, effect of rosiglitazone, 1236-1242
 - plasma, *see* Plasma homocysteine
 - serum, *see* Serum homocysteine
 - see also* S-Adenosylhomocysteine;
 - Hyperhomocysteinemia; Hypohomocysteinemia
- Homogentisic acid (HGA), urinary levels in alkaptonuria patients, effect of nitisinone, 719-728
- Hormone replacement therapy (HRT)
- and postmenopause, effect on serum adiponectin levels, 1610-1614
 - types of, effect on Lp(a), Hcy, and CRP levels, in PM women, 72-78
- Hormones, and nonesterified fatty acids, lactate, glycerol and triglycerides after 2-day VLCD, 709
- HRT, *see* Hormone replacement therapy
- HTN, *see* Hypertension
- Human arterial smooth muscle cells, effects of insulin, glucose, IGF-1, or growth hormone on formation and function of HA receptor CD44 on, 287-295
- Human chorionic gonadotropin (hCG), for adolescent boys with CDGP versus those with β -thalassemia major, 15-23
- Human growth hormone (hGH)
- correlation to plasma leptin levels, in children with short stature administered with insulin, 862-865
 - and insulin, glucose, or IGF-1, effects on formation and function of HA receptor CD44, on human arterial smooth muscle cells, 287-295
- Human immunodeficiency virus (HIV) infection, with lipodystrophy, hepatic insulin extraction and posthepatic insulin clearance rate in proportion to insulin resistance in, 171-179
- Human liver cell lines, mechanism of action of ethanol in down-regulation of 2,6-ST mRNA in, 729-734
- Hunger, and satiety, effect of pioglitazone on, 24-32
- Hx, *see* Hypoxanthine
- HX531, effect on leptin resistance without increasing plasma leptin level, in KK-A^y mice under normal dietary conditions, 573-578
- Hyaluronan receptor CD44, formation and function of, effects of insulin, glucose, IGF-1, or growth hormone, on human arterial smooth muscle cells, 287-295
- Hydrochlorothiazide, in targeting CV risk in nondiabetic patients with MS, 1065-1074
- Hydrocortisone, *see* Cortisol
- 3-Hydroxy-butylate (3-HB), plasma levels of, effects of HX531 in KK-A^y mice, 574-576
- Hydroxycobalamin, versus cyanocobalamin, effects on plasma Hcy concentrations in ESRD, 1362-1367
- Hydroxymethyl glutaryl coenzyme A (HMG-CoA) reductase inhibitors, effect on transactivation of human serum *PON1* gene, 142-150
- 1- α -Hydroxyvitamin D₃, versus alendronate, effects on BMD and bone metabolic markers, in PM asthmatic women treated with inhaled corticosteroids, 85-90
- Hyperadiponectinemia, association with thyroid hormone levels in Basedow disease, 1461-1466
- Hyperandrogenism, and menstrual function in female hypoleptinemic lipodystrophic patients, long-term effect of recombinant methionyl human leptin therapy, 255-263
- Hypercapnia, response of protein synthesis to, in rats, 841-847
- Hypercholesterolemia (HC)
- effect of SCH 58235 and high plant sterol-enriched diet on plant sterol absorption and plasma concentrations in WKY rats with, 38-48
 - primary, effects of raloxifene and low-dose simvastatin coadministration on serum lipids in PM women with, 939-946
 - relation to adhesion of monocytes to cultured endothelium in nondiabetic subjects, 1467-1471
 - sensitivity of angiotensin II-induced oxidative burst to fluvastatin in neutrophils of patients with, 1147-1154
 - in Taiwan, caused by mutations in *LDL receptor* gene, 1082-1086
- Hyperglycemia
- after androgen-deprivation therapy for prostate cancer, pioglitazone for, 55-59
 - and hepatic glucose production, effects of ASO targeted against GCCR in DM mice, 848-855
 - in poorly controlled DM, effect on brain glucose concentrations as measured by high-field magnetic resonance spectroscopy, 1008-1013

- Hyperglycemia (continued)**
postprandial, *see* Postprandial hyperglycemia
prevention in Zucker diabetic fatty rats, and gene expression in insulin-sensitive tissues determined by high-density oligonucleotide microarray analysis, effects of exercise training on, 1571-1581
- Hyperhomocysteinemia**, experimental, relationship between *S*-adenosylmethionine, *S*-adenosylhomocysteine and asymmetric dimethylarginine, in healthy human subject during, 351-360
- Hyperinsulinemia**
contributions of oxidative and nonoxidative glucose disposal to, in elderly versus younger men with similar and smaller BMI and waist circumferences, 751-752
predicts hepatic fat content in healthy individuals with normal ALT concentrations, 1566-1570
prolonged, blood pressure changes and plasma catecholamine levels during, 391-396
protein anabolic response to, influence of sex, 1529-1535
- Hyperleptinemia**, glucocorticoid-induced, energy consumption in, 876-880
- Hyperlipidemia**
diet-induced insulin-resistant, association of increased diacylglycerol acyltransferase activity with triglyceride accumulation in tissues of, 403-409
NIDDM complicated by, effect of low-dose atorvastatin on circulating MCP-1, 1225-1229
types IIa and IIb, comparative study of beneficial effects of pravastatin, atorvastatin and fenofibrate in, 677-681
see also Hypercholesterolemia
- Hypertension (HTN)**
and albuminuria, association with circulating APAAs in NIDDM, 188-193
association with cellular insulin resistance in EBV-transformed lymphoblasts from young Japanese men, 370-375
effect of selective angiotensin II receptor antagonism on whole-body insulin sensitivity and muscle glucose transport in, 1659-1668
and insulin resistance, fructose-induced, functional interaction of AT1R and AT2R in, 157-164
and NIDDM, effect of rosiglitazone on novel atherosclerotic risk factors in, 1236-1242
and obesity and IDDM, levels of sICAM-1, sVCAM-1 and selectins in children and adolescents with, 1020-1026
- Hypertrophy**, of RV, and hypoxia-induced anorexia, role in lactate transport and monocarboxylate transporter expression in rat muscle, 634-644
- Hyperuricemia**, and alcohol drinking, possible interactions with eNOS among Japanese, 1302-1308
- Hypoadiponectinemia**
association with arteriosclerosis obliterans and IHD, 653-656
caused by chronic blockade of nitric oxide synthesis, 482-487
- Hypoalbuminemia**, CRP levels and prevalence of chronic infections in, 33-37
- Hypofibrinolysis**, and thrombophilia in women with miscarriages and RPL, 1346-1347
- Hypoglycemia**
adrenal or lumbar SNA induced by, effect of captopril on reflex increases, 1679-1686
recurrent, effect of combined CRH and glucocorticoid deficiency on counterregulatory responses after, 1259-1265
- Hypogonadism**, in adolescent boys with β -thalassemia major, effects of hCG therapy on, 20-22
- Hypohomocysteinemia**, experimental, relationship between *S*-adenosylmethionine, *S*-adenosylhomocysteine and asymmetric dimethylarginine, in healthy human subject during, 351-360
- Hypoleptinemic lipodystrophy**, hyperandrogenism, menstrual function, and pituitary function in patients, long-term effect of recombinant methionyl human leptin, 255-263
- Hypothalamic neuropeptides**, role in neurophysiologic modulation of obesity, 1207-1209
- Hypothermia**, effects on tissue protein synthesis in rats, 841-847
- Hypothyroidism**, methimazole-induced, effect on homocysteine concentrations, 460-465
- Hypoxanthine (Hx)**, urinary excretion and plasma concentrations, influence of allopurinol on urinary purine loss after repeated sprint exercise in man, 1271-1273
- Hypoxia**, anorexia induced by, and RV hypertrophy, role in lactate transport and monocarboxylate transporter expression in rat muscle, 634-644
- IAGTT**, *see* Intra-arterial glucose tolerance test
- Ibuprofen**, NO-1886, effect on expression of UCP3 in skeletal muscle and fat accumulation in HF diet-induced obesity, 1587-1592
- ICAM-1 gene**, *see* Intercellular adhesion molecule-1 gene
- IDDM**, *see* Insulin-dependent diabetes mellitus
- Idiopathic short stature (ISS)**, leptin responses to insulin administration in children with, 862-865
- IGF-1**, *see* Insulin-like growth factor-1
- IGFBP-1**, *see* Insulin-like growth factor-binding protein 1
- IGF-BPs**, *see* Insulin-like growth factor-binding protein(s)
- IHD**, *see* Ischemic heart disease
- IL-6**, *see* Interleukin-6
- IL-8**, *see* Interleukin-8
- IMCL**, *see* Intramyocellular lipid(s)
- IMGU**, *see* Insulin-mediated glucose uptake

- IMT (intimal-media thickness), *see* Carotid artery
intimal-media thickness (IMT)
- Inflammation
and activation, endothelial, in prepubertal obese Turkish children, 1384-1389
and AT, role in neurophysiologic modulation of obesity, 1204-1205
- Inflammatory cytokines, effects of diet and/or exercise in postmenopausal NIDDM women, 866-875
- Inflammatory markers, and insulin sensitivity in overweight and obese girls, effects of aerobic exercise training, 1472-1479
- Inflammatory responses, in NIDDM with CAD after coronary angioplasty, effect of rosiglitazone on, 590-597
- Inhaled corticosteroids, effects of alendronate on BMD and bone metabolic markers in PM asthmatic women treated with, 85-90
- Inosine, urinary excretion and plasma concentrations, influence of allopurinol on urinary purine loss after repeated sprint exercise in man, 1271-1273
- Insulin
concentration of, and FPG after 2-day VLCD, 708
continually high levels, effect on Akt phosphorylation and glucose transport in human myoblasts, 1687-1693
effect on VEGF, IL-8, PAI-1, and IL-6 release by human adipocytes, 220-226
endothelial activation and inflammation in prepubertal obese Turkish children, 1384-1389
and glucose, IGF-1, or growth hormone, effects on formation and function of HA receptor CD44, on human arterial smooth muscle cells, 287-295
glucose transport stimulated by, in human omental versus SC adipocytes, 781-785
hepatic, *see* Hepatic insulin
long-term and immediate effects of intranasal administration on BP and MSNA, 1356-1361
and plasma glucose levels during HC induction of hyperleptinemia, 877-878
plasma leptin responses to, in children with short stature, 862-865
protein anabolic response to, influence of sex, 1529-1535
role in neurophysiologic modulation of obesity, 1203
serum, *see* Serum insulin
STZ-DM fasted rats treated with, effect of impaired glucose uptake on postexercise glycogen repletion in SM, 1420-1427
subcutaneous versus buccal spray, in IDDM patients, 930-934
withdrawal of, and VLCD, effect on endogenous glucose production in obese NIDDM, 705-712
see also Diabetes mellitus; Gestational diabetes mellitus; Hyperinsulinemia; Hypoglycemia; Insulin-dependent diabetes mellitus; Insulinemia; Insulin-like growth factor-1; Insulin-mediated
Insulin (continued)
glucose uptake; Insulin release and secretion; Insulin resistance; Insulin sensitivity; Insulin signaling; Non-insulin-dependent diabetes mellitus; Streptozotocin-induced diabetes mellitus
Insulin action, in response to endurance training, association with PPAR γ 2 Pro12Ala genotype, in men and women, 97-102
Insulin-dependent diabetes mellitus (IDDM)
association with functional polymorphism of Z-DNA-forming motif of promoter of *SLC11A1* gene in Japanese subjects, 628-633
biokinetics of buccal spray insulin in, 930-934
and obesity and hypertension, levels of sICAM-1, sVCAM-1 and selectins in children and adolescents with, 1020-1026
Insulinemia, or sulfonylurea usage, role in weight gain in NIDDM patients, 28-29
Insulin-like growth factor-1 (IGF-1)
correlation to plasma leptin levels, in children with short stature administered with insulin, 862-865
and GH, chronic excess, cardiac and metabolic effects in young adults with pituitary gigantism, 1174-1180
and IGF-BPs, circulating, genetic regulation of variation in human pedigrees, 975-981
and insulin, glucose, or growth hormone, effects on formation and function of HA receptor CD44, on human arterial smooth muscle cells, 287-295
and insulin sensitivity in overweight and obese girls, effects of aerobic exercise training, 1472-1479
Insulin-like growth factor-binding protein(s) (IGF-BPs), and IGF-1, circulating, genetic regulation of variation in human pedigrees, 975-981
Insulin-like growth factor-binding protein 1 (IGFBP-1), altered response to nutritional deprivation in NIDDM, 275-280
Insulin-mediated glucose uptake (IMGU), relation to excess adiposity as determined by BMI and WC, 1323-1328
Insulin promoter factor-1 (IPF-1) gene, contribution of missense mutation to MODY4 in Italian family with NIDDM and GDM, 983-988
Insulin receptor, effect of continually high insulin levels on Akt phosphorylation and glucose transport in human myoblasts, 1687-1693
Insulin receptor activation, and glucose transport, insulin-stimulated, analysis in cultured skeletal muscle cells from obese subjects, 598-603
Insulin receptor substrate (IRS), effect of continually high insulin levels on Akt phosphorylation and glucose transport in human myoblasts, 1687-1693
Insulin release and secretion
association with fasting serum nonesterified fatty acids, effect of westernized lifestyle in Japanese men, 713-718

- Insulin release and secretion (continued)
and sensitivity, relationship to MS at follow-up in women with and without GDM in index pregnancy, 1115–1121
short-term effects of FAs as influenced by their chain length, spatial configuration and degree of unsaturation, 1329–1336
- Insulin Resistance Atherosclerosis Study, 264–270
- Insulin resistance (IR)
and adiposity, influence on lipoprotein size and subclass concentrations, 264–270
and apo C-III bound to apo B-containing lipoproteins, in Cherokee Indian youth, 180–187
association with C161→T polymorphism in PPAR γ in Hispanic and non-Hispanic white women, 1552–1556
cellular, in Epstein-Barr virus-transformed lymphoblasts from young insulin-resistant Japanese men, 370–375
and C-reactive protein, reciprocal association with adipocytokines, in clinically healthy men, 439–444
effect on intestinal assembly and secretion of apo B48-containing Lp particles in fasting state, 689–697
effects of metformin on, 235–239
and glucose uptake, effect of p38 MAP kinase inhibition in 3T3-L1 adipocytes, 895–901
in HIV infection with lipodystrophy, hepatic insulin extraction and posthepatic insulin clearance rate in proportion to, 171–179
and hypertension, fructose-induced, functional interaction of AT1R and AT2R in, 157–164
metformin-diet benefits in women with PCOS and, 113–121
model of, metabolic responses with ET antagonism in, 735–740
in MS, ultrasonography in prediction of, 1230–1235
in NIDDM, *see* Insulin resistance (IR), in NIDDM
and normal TGs, relation to PH-LPL activity in African Americans, 902–909
in obesity, *see* Insulin resistance (IR), in obesity
and plasma resistin, effects of FFAs in awake rats, 1142–1146
and postprandial TGs in primary renal disease, 821–828
predicts hepatic fat content in healthy individuals with normal ALT concentrations, 1566–1570
progression in nondiabetic Zucker rats, age and muscle-type modulated role of intramyocellular lipids in, 522–528
and serum insulin, glucose response to IVGTT and pancreatic insulin content, effects of camostat, 623
see also Homeostasis model assessment insulin resistance (HOMA_{IR}) index
- Insulin resistance (IR), in NIDDM
and cardiovascular autonomic dysfunction, association with high-sensitivity C-reactive protein, 552–558
effects of red wine consumption on endothelial function and, 306–313
- Insulin resistance (IR), in NIDDM (continued)
non-obese Japanese subjects
relationship to soluble E-selectin, triglycerides, and leptin, 376–380
relation to TNF- α activity, 1297–1301
parameters of, correlation with adiponectin-leptin ratio, 281–286
- Insulin resistance (IR), in obesity
analysis of insulin receptor activation and glucose transport in skeletal muscle cells, 598–603
relation to high-sensitive CRP, TNF- α , blood pressure and lipids before and after weight loss in children, 1155–1161
- Insulin-resistant hyperlipidemia, diet-induced, association of increased diacylglycerol acyltransferase activity with triglyceride accumulation in tissues of, 403–409
- Insulin secretion, *see* Insulin release and secretion
- Insulin secretory capacity, contribution of *GPR40* gene Arg211His polymorphism on variation of, in Japanese men, 296–299
- Insulin secretory defects, and dyslipidemia and histological abnormalities of pancreas, effects of synthetic PI camostat in genetically obese and diabetic rats, 619–627
- Insulin sensitivity (*S_i*)
and adipocyte plasma membrane fatty acid composition, effects of dietary *trans*-FA on, 240–246
and adipocytokine levels, effect of pravastatin in healthy nondiabetic patients, 947–951
and body weight, body fat, adiponectin and inflammatory markers in overweight and obese girls, effects of aerobic exercise training, 1472–1479
and body weight control, effect of RVF containing high cyclo (his-pro), 1480–1489
effect of short-term leptin treatment on FA uptake and oxidation in muscle of high fat-fed rats, 1218–1224
hepatic, associated with insulin signaling, effect of ET antagonism in Zucker fatty rats, 1515–1523
longitudinal measures of, effects of *PPAR* γ 2 Pro12Ala polymorphism in African Americans and whites, 910–917
in offspring rats, effects of high saturated fat versus high-CHO diet during pregnancy and lactation, 1316–1322
postprandial, modifying effect of vagal nerve stimulation on, 579–583
and secretion, relationship to MS at follow-up in women with and without GDM in index pregnancy, 1115–1121
SM, effect of 2-day VLED in obese NIDDM patients on insulin therapy, 1669–1678
spectrum of, in Korean population, 1644–1651
in transgenic rabbits, effect of high LPL activity on, 132–138
whole-body, *see* Whole-body insulin sensitivity (*S_i*)

- Insulin sensitivity indices (ISI), of glucose and free fatty acid metabolism, in relation to serum lipids, in obese children and adolescents, 397-402
- Insulin signaling
- hepatic insulin sensitivity associated with, effect of ET antagonism in Zucker fatty rats, 1515-1523
 - PI3K, inhibition leads to decreased insulin-stimulated adiponectin secretion from 3T3-L1 adipocytes, 1636-1643
- Insulin signaling proteins, in male offspring of high-fat fed rats, 500-507
- Insulin therapy, effect of 2-day VLED on SM insulin sensitivity in obese NIDDM patients on, 1669-1678
- Insulin tolerance test (ITT), plasma leptin responses to, in children with short stature, 862-865
- Intercellular adhesion molecule-1 (ICAM-1), *see* Soluble intercellular adhesion molecule-1
- Intercellular adhesion molecule-1 (ICAM-1) gene, K469E polymorphism of, association with plasma fibrinogen level in NIDDM, 381-386
- Interleukin-6 (IL-6)
- and insulin sensitivity in overweight and obese girls, effects of aerobic exercise training, 1472-1479
 - release by human adipocytes, effect of insulin on, 220-226
 - serum, *see* Serum interleukin-6 (IL-6)
- Interleukin-8 (IL-8), release by human adipocytes, effect of insulin on, 220-226
- Intestinal absorption, in hypercholesterolemic wild-type Kyoto rats, effect of SCH 58235 and high plant sterol-enriched diet on, 38-48
- Intimal hyperplasia, and serum Hcy and smooth muscle proliferation, effect of rosiglitazone in high methionine-fed Sprague-Dawley rats, 645-652
- Intimal-media thickness (IMT), *see* Carotid artery intimal-media thickness (IMT)
- Intra-arterial glucose tolerance test (IAGTT), effects of soy-rich diet and stevioside in NIDDM rats, 1185
- Intracellular triglycerides (TGs) content, in skeletal muscle and liver, inverse correlation with serum adiponectin concentration, in Japanese NIDDM subjects, 775-780
- Intramyocellular lipid(s) (IMCL), age and muscle-type modulated role in insulin resistance progression, in nondiabetic Zucker rats, 522-528
- Intranasal administration, insulin, long-term and immediate effects on BP and MSNA, 1356-1361
- Intrarenal blood flow, and role of NO in diabetic rats, 788-792
- Intravenous glucose tolerance test (IVGTT), glucose response to, and serum insulin, IR and pancreatic insulin content, effects of camostat, 623
- In vitro neutrophil oxidative burst, effects of propylthiouracil and thiamazole, 300-305
- IPF-1 gene, *see* Insulin promoter factor-1 gene
- IR, *see* Insulin resistance
- Iron loading, effect on cholesterol accumulation and macrophage scavenger receptor I expression in THP-1 mononuclear phagocytes, 453-459
- IRS, *see* Insulin receptor substrate
- Ischemic heart disease (IHD), and arteriosclerosis obliterans, association with hypoadiponectinemia, 653-656
- ISI, *see* Insulin sensitivity indices
- Isoenergetic protein ingestion, versus CHO supplementation, combined with resistance training, effect on muscle fiber size and muscle strength, 151-156
- Isotopomer spectral analysis, of cholesterol synthesis intermediates, in patients with cerebrotendinous xanthomatosis, 335-344
- ISS, *see* Idiopathic short stature
- Italian subjects, with NIDDM
- and GDM, IPF-1/MODY4 gene missense mutation, 983-988
 - genetic linkage of glucagon receptor gene, 786-787
- ITT, *see* Insulin tolerance test
- IVGTT, *see* Intravenous glucose tolerance test
- Japanese Americans, *see* Race and ethnicity
- Japanese NIDDM subjects
- association of microalbuminuria with G1704T and G82S polymorphisms of *RAGE* gene, 488-491
 - inverse correlation between serum adiponectin concentration and HLC, 775-780
 - men, association between serum DHEAS levels and NIDDM progression, 669-676
 - non-obese
 - soluble E-selectin, leptin, triglycerides, and insulin resistance, 376-380
 - three measures of TNF- α and insulin resistance, 1297-1301
- Japanese subjects
- association between functional polymorphism of Z-DNA-forming motif of promoter of *SLC11A1* gene and IDDM, 628-633
 - men
 - cellular insulin resistance in Epstein-Barr virus-transformed lymphoblasts obtained from, 370-375
 - contribution of *GPR40* gene Arg211His polymorphism on variation of insulin secretory capacity, 296-299
 - effect of westernized lifestyle on the association between fasting serum nonesterified fatty acids and insulin secretion in, 713-718
- NIDDM, *see* Japanese NIDDM subjects
- nonobese, effects of thorough mastication on postprandial plasma glucose concentrations, 1593-1599
 - possible interactions of eNOS genotype with alcohol drinking and walking time for high serum uric acid levels among, 1302-1308

- JNK activation, mediators of, hepatic stress in response to fructose metabolism, 1194
- K469E polymorphism, of *ICAM-1* gene, association with plasma fibrinogen level in NIDDM, 381-386
- Ketone bodies, metabolism in lean versus obese women, 1542-1545
- Ketosis, transient, classic high-fat ketogenic diet induces marked changes in fatty acid metabolism in rats in absence of, 1127-1132
- Kidney function, effects of chronic chromium picolinate treatment in uninephrectomized rats, 1243-1249
- Korea, limitation of the validity of HOMA of insulin resistance in, 206-211
- Korean subjects
 apparently healthy men, relationship between normal WBC count and cardiorespiratory fitness, 1448-1452
 spectrum of insulin sensitivity in, 1644-1651
 women
 association of circulating OPG with age, WHR, serum TC, and LDL-C levels in, 49-54
 postmenopausal, association among serum ferritin, ALT levels and MS in, 1510-1514
- LA, *see* Alpha (α)-lipoic acid
- Lactate, *see* Blood lactate; Lactic acid
- Lactate dehydrogenase isoforms, and monocarboxylate transporter expression, role of hypoxia-induced anorexia and RV hypertrophy in rat muscle, 634-644
- Lactate transport
 during exercise in NIDDM postmenopausal women, 1105
 and monocarboxylate transporter expression, role of hypoxia-induced anorexia and RV hypertrophy in rat muscle, 634-644
- Lactation, and pregnancy, high saturated fat versus high CHO diet during, effects on insulin sensitivity in offspring rats, 1316-1322
- Lactic acid, and nonesterified fatty acids, glycerol, triglycerides and hormones after 2-day VLCD, 709
- Lanreotide Autogel, pharmacokinetics and population pharmacodynamic analysis, 1276-1281
- LBW, *see* Low birth weight
- LDL, *see* Low-density lipoprotein(s)
- LDL-C, *see* Low-density lipoprotein-cholesterol
- Lean subjects, *see* entries beginning with terms:
 Non-obese subjects
- Leptin
 association with urinary isoprostane in Japanese Americans, 194-199
 and circulating IGFs, genetic regulation of variation in human pedigrees, 975-981
 effect on lipogenic enzyme gene expression through modification of SREBP-1c gene expression in WAT of aging rats, 1041-1047
 effect of pravastatin in healthy nondiabetic patients, 947-951
- Leptin (continued)
 long-term effects of recombinant human leptin on hyperandrogenism, menstrual function, and pituitary function in hypoleptinemic lipodystrophic patients, 255-263
 plasma, *see* Plasma leptin
 and plasma cortisol levels, during HC induction of hyperleptinemia, 878
 relationship to soluble E-selectin, triglycerides, and insulin resistance, in nonobese Japanese with NIDDM, 376-380
 role in neurophysiologic modulation of obesity, 1205-1206
 short-term treatment, effect on FA uptake and oxidation in muscle of high fat-fed rats, 1218-1224
 sTNF-R, BMI, adiponectin and serum lipid profile in nonobese Japanese NIDDM patients, 1297-1301
 TNF- α -associated expression of, in dimethylnitrosamine and bile duct ligation-induced liver cirrhosis, 445-452
- Leptin-adiponectin ratio, correlation with parameters of insulin resistance, in NIDDM, 281-286
- Lifestyle
 changes, effect on insulin action in obese African Americans, 960-965
 westernized, effect on the association between fasting serum nonesterified fatty acids and insulin secretion in Japanese men, 713-718
- Linoleic acid, effects on vascular reactivity in healthy adult rats, 1600-1609
- α -Linolenic acid, systematic review and meta-analysis of impact on selected arrhythmia outcomes in animal models, 1557-1565
- Lipemic responses, to HCHF diet, LDL apo B production in laboratory opossums exhibiting high and low, 1075-1081
- Lipid(s)
 blood, *see* Blood lipids
 elevated concentrations of, effect on human SM gene expression, 952-959
 and fatty acids, hepatic, physiological relevance of apoE recycling in primary mouse hepatocytes, 1311-1312
 and glucose, comparison of immediate response to meals enriched with *cis*- or *trans*-FAs in overweight individuals with differing FABP2 genotype, 1652-1658
 and glucose metabolism after liver transplantation in inbred rats, 881-890
 plasma, *see* Plasma lipids
 relation to high-sensitive CRP, TNF- α , blood pressure and insulin resistance before and after weight loss in obese children, 1155-1161
 serum, *see* Serum lipids
 see also entries beginning with element: Lip- and specific lipids
- Lipid disorders clinic patients, Hcy, glycine betaine, and *N,N*-dimethylglycine in, 1-14

- Lipid-lipoprotein profile, according to apo E genotype, 1016
- Lipid-lowering drugs, comparative study of beneficial effects of pravastatin, atorvastatin and fenofibrate in types IIa and IIb hyperlipidemia patients, 677-681
- Lipid metabolism, liver, of male offspring of high-fat fed rats, 500-507
- Lipid oxidation rates
and glucose after 2-day VLCD, 709
in whole body lipolysis, 496
- Lipid profile
beneficial effects of pravastatin, atorvastatin and fenofibrate in types IIa and IIb hyperlipidemia patients, 677-681
and carotid artery IMT, in adults with growth hormone deficiency after long-term growth hormone replacement, 321-329
relationship to soluble E-selectin, leptin, and insulin resistance, in nonobese Japanese with NIDDM, 376-380
serum, *see* Serum lipid profile
- Lipid synthesis, and insulin sensitivity in offspring rats, effects of high saturated versus high-CHO diet during pregnancy and lactation, 1319
- Lipodystrophy
HIV infection with, hepatic insulin extraction and posthepatic insulin clearance rate in proportion to insulin resistance in, 171-179
hypoleptinemic, hyperandrogenism, menstrual function, and pituitary function in patients, long-term effect of recombinant methionyl human leptin, 255-263
- Lipogenic enzyme, gene expression, effect of leptin through modification of SREBP-1c gene expression in WAT of aging rats, 1041-1047
- α (alpha)-Lipoic acid (LA), supplementation with, effects on exercise-induced blood coagulation activation, 815-820
- Lipolysis, whole body, resting and terbutaline-stimulated, effect of sex and β_2 adrenergic receptor haplotype on, 492-499
- Lipolytic pathways, dependent on ANP and adrenergic receptor, differential regulation in human SAT, 122-131
- Lipoprotein (Lp)
atherogenic subfraction profile in preeclamptic women with and without high TGs, 1504-1509
concentration, particle size, and subclass concentration, influence of insulin resistance and adiposity on, 264-270
containing apo B, and insulin resistance in Cherokee Indian youth, 180-187
containing apo B48, intestinal assembly and secretion in fasting state, 689-697
effects of raloxifene and low-dose simvastatin coadministration in PM women with primary hypercholesterolemia, 939-946
- Lipoprotein (Lp) (continued)
influence on whole-blood viscosity at multiple shear rates, 764-768
see also High-density lipoprotein; Lipoprotein(a); Lipoprotein lipase; Low-density lipoprotein
- Lipoprotein(a) [Lp(a)]
and baseline LDL-C and *Apo*II polymorphism of *LDL-receptor* gene, influence on LDL-C response to pravastatin, 741-747
levels in PM women, effects of types of HRT, 72-78
in patients with NIDDM and hypertension, effect of rosiglitazone, 1236-1242
plasma, *see* Plasma lipoprotein(a)
- Lipoprotein lipase (LPL)
complete deletion, effects of DAG administration on serum TAGs in patient homozygous for, 67-71
high activity of, effect on insulin sensitivity in transgenic rabbits, 132-138
postheparin, *see* Postheparin-lipoprotein lipase
preheparin serum, *see* Preheparin serum lipoprotein lipase (LPL)
S447X variant, effect on postprandial apo B-48 clearance, 1499-1503
- Lipoprotein lipase (LPL) activator, NO-1886 (ibrolipim), effect on expression of UCP3 in skeletal muscle and fat accumulation in HF diet-induced obesity, 1587-1592
- Lipoprotein (Lp)-lipid profile, serum, and CHO intake and nutrient intake ratios, effects of beef versus soy-based food supplements in older men who perform RT, 769-774
- Liver
adaptations to sucrose and fructose, 1189-1201
effects of chronic ethanol feeding on activities and RSRs of various sialidases in, 1056-1064
gene expression in, and hyperglycemia prevention in Zucker diabetic fatty rats, effects of exercise training on, 1571-1581
transgenic expression of mutant PPAR γ in, effects on fasting-induced and HF diet-induced steatosis in mice, 1490-1498
see also entries beginning with *element*: Hepat- and *specific conditions*
- Liver cirrhosis, dimethylnitrosamine and bile duct ligation-induced, TNF- α -associated leptin expression in, 445-452
- Liver function, improvement in NIDDM patients treated with thiazolidinediones, 529-532
- Liver lipid metabolism, of male offspring of high-fat fed rats, 500-507
- Liver transplantation, glucose and lipid metabolism after, in inbred rats, 881-890
- Liver weight, effects of treatment of ASO against GCCR, 853
- LMS, *see* Lysosomal membrane sialidase
- L-NAME, *see* L- ω -Nitro-L-arginine methyl ester

- Long-term administration, of synthetic PI camostat, effects on dyslipidemia, insulin secretory defects and histological abnormalities of pancreas in genetically obese and diabetic rats, 619-627
- Long-term effects
- of alcohol administration on synthesis of myofibrillar and sarcoplasmic proteins in heart, 212-219
 - of early exposure to GH and IGF-1 excess on cardiovascular and metabolic parameters in young adults with pituitary gigantism, 1174-1180
 - and immediate effects of intranasal administration of insulin on BP and MSNA, 1356-1361
 - of recombinant methionyl human leptin therapy on hyperandrogenism, menstrual function, and pituitary function in hypoleptinemic lipodystrophic patients, 255-263
- Long-term growth hormone replacement, carotid artery IMT and lipid profile in growth hormone deficiency after, 321-329
- Low back pain, as somatic marker of depression, S1:41-42
- Low birth weight (LBW), susceptibility to diabetogenic effects of prolonged FFA exposure in young men with, 1398-1406
- Low-density lipoprotein(s) (β -lipoprotein; LDL)
- apo B production in laboratory opossums exhibiting high and low lipemic responses to dietary cholesterol and fat, 1075-1081
 - atherogenicity of, effect of weight loss associated with reduced intake of CHO in premenopausal women, 1133-1141
 - metabolism, effect of iron loading in THP-1 mononuclear phagocytes, 453-459
 - oxidation, and plasma antioxidant capacity in men after short-term cranberry juice consumption, 856-861
 - size and subclasses of, as markers of clinically apparent and non-apparent atherosclerosis in NIDDM, 227-234
 - uptake in human lymphocytes, influence of cyclosporine, 1620-1625
- Low-density lipoprotein-cholesterol (β -lipoprotein-cholesterol; LDL-C)
- association with circulating OPG in healthy Korean women, 49-54
 - concentrations, influence of apo E genotype on reliability of Friedewald formula in estimation of, 1014-1019
 - response to pravastatin, influence of baseline LDL-C, Lp(a) and *AvalI* polymorphism of *LDL-receptor* gene, 741-747
 - serum, *see* Serum low-density lipoprotein-cholesterol
- Low-density lipoprotein (LDL) receptor gene
- AvalI* polymorphism of, and baseline LDL-C and Lp(a), influence on LDL-C response to pravastatin, 741-747
 - mutations of, as cause of hypercholesterolemia in Taiwan, 1082-1086
- Low-dose atorvastatin, effect on circulating MCP-1 in patients with NIDDM complicated by hyperlipidemia, 1225-1229
- Lp, *see* Lipoprotein
- Lp(a), *see* Lipoprotein(a)
- LPL, *see* Lipoprotein lipase
- L-selectin, *see* Soluble L-selectin
- Lumbar sympathetic nerve activity (SNA), hypoglycemia-induced, effect of captopril on reflex increases in, 1679-1686
- Lung, fibrinogen metabolism in, 271-274
- Lymphoblasts, Epstein-Barr virus-transformed, cellular insulin resistance in, 370-375
- Lymphocytes, human, influence of cyclosporine on LDL uptake in, 1620-1625
- Lysosomal membrane sialidase (LMS), activities and RSRs in rat liver, effects of chronic ethanol feeding, 1056-1064
- Macronutrient consumption, during HC induction of hyperleptinemia, 878
- Macrophage scavenger receptor I, expression of, and cholesterol accumulation, effect of iron loading in THP-1 mononuclear phagocytes, 453-459
- Magnetic resonance spectroscopy, high-field, measurement of brain glucose concentrations in poorly controlled DM, 1008-1013
- Major depressive disorder (MDD; unipolar depression)
- abdominal AT and serum levels of prothrombic factors in younger, premenopausal women with, 918-924
 - genetic influences on susceptibility to, S1:16-19
 - genetics and genomics, S1:11
 - linkage analysis, S1:11-12
 - overlap with bipolar depression, S1:11
- Malaria, glucose production and plasma adiponectin in, 60-66
- Malnutrition, and nonthyroidal illness syndrome after stroke, 699-704
- Malondialdehyde (MDA), plasma levels, effects of α -lipoic acid, 817-818
- Massive tendon xanthomatosis, familial, with decreased HDL-mediated cholesterol efflux, 1095-1101
- Mastication, thorough, effects on postprandial plasma glucose concentrations in nonobese Japanese subjects, 1593-1599
- Maturity-onset diabetes of the young 4 (MODY4), contribution of *IPF-1* gene missense mutation to, in Italian family with NIDDM and GDM, 983-988
- MCP-1, *see* Monocyte chemoattractant protein-1
- MCR, *see* Metabolic clearance rate
- MCTs, *see* Medium chain triacylglycerols
- MDA, *see* Malondialdehyde
- MDD, *see* Major depressive disorder
- Meals, *see* Diet
- Medical illness, association with psychiatric illness, S1:53-54

- Medium chain triacylglycerols (MCTs), effects of dietary DAG on, in patient with LPL deletion, 67-71
- Melanin-concentrating hormone, role in neurophysiologic modulation of obesity, 1208
- Men
- apparently healthy Korean, *see* Korean subjects
 - contribution of age and declining androgen levels to features of MS in, 1034-1040
 - Japanese, *see* Japanese NIDDM subjects; Japanese subjects
 - middle-aged and young athletes versus sedentary, effects of exercise at moderate and hard intensity on substrate oxidation, 1411-1419
 - with NIDDM and impaired glucose tolerance, impact of glucagon response on postprandial hyperglycemia, 1168-1173
 - plasma antioxidant capacity and OxLDL levels in, after cranberry juice consumption, 856-861
 - and women, healthy aged, effects of GH and/or sex steroids on whole-body protein turnover, 1162-1167 *see also* Sex
- Menopause
- and depression, S1:51
 - and HRT, effect on serum adiponectin levels, 1610-1614
- Menstrual function, and hyperandrogenism, in female hypoleptinemic lipodystrophic patients, long-term effect of recombinant methionyl human leptin therapy, 255-263
- Mental illness, association with somatic disorders, S1:53-54
- Mesenteric adipose tissue, accumulation, and UCP3 expression in skeletal muscle, effect of NO-1886 on HF diet-induced obesity, 1587-1592
- Mesenteric fatty tissue weight, effects of HX531 in KK-A^y mice, 574-576
- Messenger ribonucleic acid (mRNA)
- adipocyte expression in thigh SAT, effect of PI-based HAART, 561-567
 - expression in insulin-sensitive tissues determined by high-density oligonucleotide microarray analysis and hyperglycemia prevention in Zucker diabetic fatty rats, effects of exercise training on, 1571-1581
 - expression of adiponectin receptor 1 in obese (fa/fa) and lean Zucker rats, 998
 - expression of lipogenic enzyme, effect of leptin through modification of SREBP-1c gene expression in WAT of aging rats, 1041-1047
 - expression of PPARs in obese (fa/fa) and lean Zucker rats, 999
 - hepatic expression of enzymes, effect of apoE recycling, 1312-1313
 - of mutant PPAR γ , effect of transgenic expression in liver on fasting-induced and HF diet-induced steatosis, 1490-1498
 - SM gene expression, effect of elevated lipid concentrations, 952-959
- Messenger ribonucleic acid (mRNA) (continued)
- UCP3 expression in skeletal muscle, and fat accumulation, effect of NO-1886 on HF diet-induced obesity, 1587-1592
- MET, *see* Metformin
- Metabolic clearance rate (MCR), and spontaneous secretion, of GH in overweight youth, pubertal alterations in, 1374-1383
- Metabolic genes, substrate availability and transcriptional regulation in human SM during recovery from exercise, 1048-1055
- Metabolic parameters
- relationship to fasting plasma ghrelin in children and adolescents, 925-929
 - relationship to serum resistin level, among healthy subjects, 471-475
- Metabolic syndrome (insulin resistance syndrome; MS; syndrome X)
- association with lower serum free thyroxine levels in Chinese population, 1524-1528
 - association with low levels of adiponectin, 476-481
 - features of, contribution of age and declining androgen levels in men with, 1034-1040
 - at follow-up in women with and without GDM in index pregnancy, 1115-1121
 - and NIDDM, preventive effects of soy-based diet supplemented with stevioside in development of, 1181-1188
 - and obesity, development into NIDDM and CV disease, sex difference in subjects with normal glucose metabolism, 800-808
 - prevalence of, contribution of oral glucose tolerance test in, 542-547
 - relation to adhesion of monocytes to cultured endothelium in nondiabetic subjects, 1467-1471
 - relation to serum ferritin and ALT levels in Korean postmenopausal women, 1510-1514
 - targeting CV risk in nondiabetic patients with, 1065-1074
 - and visceral fat, ultrasonography for evaluation of, 1230-1235
- Metformin (dimethylguanidine; Glucophage; MET)
- and diet, benefits in women with PCOS and IR, 113-121
 - effect on intracellular production of ROS in aortic ECs, 829-834
 - effects on glucose tolerance, insulin resistance, beta cell function, and sTNFR levels, 235-239
 - and rosiglitazone, effects on plasma concentrations of resistin, in NIDDM, 314-320
 - in targeting CV risk in nondiabetic patients with MS, 1065-1074
- Methimazole, hypothyroidism induced by, effect on homocysteine concentrations, 460-465
- Metronidazole, effect on serum lipids and colonic microflora, 103-112
- MI, *see* Myocardial infarction

- Microalbuminuria, association with G1704T and G82S polymorphisms of *RAGE* gene, in Japanese NIDDM patients, 488-491
- Middle-aged men
father's BMI, associations with 18-year-old son's waist circumferences, birth weight, and serum hepatic enzyme levels, 466-470
and young athletes versus sedentary men, effects of exercise at moderate and hard intensity on substrate oxidation, 1411-1419
- Middle-aged women, mother's BMI, associations with 18-year-old son's waist circumferences, birth weight, and serum hepatic enzyme levels, 466-470
- Miscarriage, sporadic, etiologies, role of *factor V Leiden* gene mutation, high factor VIII and high PAI activity, 1345-1349
- Missense mutation, of *IPF-1* gene, contribution to MODY4 in Italian family with NIDDM and GDM, 983-988
- MODY4, *see* Maturity-onset diabetes of the young 4
- Monoamines, neurobiological basis of depression, S1:25-26
- Monocarboxylate transporter, expression, and lactate transport, role of hypoxia-induced anorexia and RV hypertrophy in rat muscle, 634-644
- Monocarboxylate transporter 1, during exercise in NIDDM postmenopausal women, 1105
- Monocyte chemoattractant protein-1 (MCP-1)
circulating, *see* Circulating monocyte chemoattractant protein-1
plasma, *see* Plasma monocyte chemoattractant protein-1
- Monocytes, adhesion to cultured endothelium, relation to hypercholesterolemia and MS in nondiabetic subjects, 1467-1471
- Mononuclear phagocytes, THP-1, effect of iron loading on cholesterol accumulation and macrophage scavenger receptor I expression in, 453-459
- Mood disorders, relation to glucocorticoids and depression, S1:20-23
- mRNA, *see* Messenger ribonucleic acid
- MS, *see* Metabolic syndrome
- MSNA, *see* Muscular sympathetic nervous activity
- Muscle, effect of short-term leptin treatment on FA uptake and oxidation in high fat-fed rats, 1218-1224
- Muscle fiber
size, effect of resistance training combined with timed protein ingestion on, 151-156
type of intramyocellular lipids, role in progression of insulin resistance in nondiabetic Zucker rats, 522-528
- Muscle glucose transport, and whole-body insulin sensitivity, effect of selective angiotensin II receptor antagonism on hypertensive TG(mREN2)27 rats, 1659-1668
- Muscle hypertrophy, effect of resistance training combined with timed protein ingestion on, 151-156
- Muscle metabolites, before and after exercise in insulin-treated STZ-DM fasted rats, 1424
- Muscle strength, effect of resistance training combined with timed protein ingestion on, 151-156
- Muscular sympathetic nervous activity (MSNA), long-term and immediate effects of intranasal administration of insulin, 1356-1361
- Mutation(s)
of *factor V Leiden* gene, and high factor VIII and high PAI activity, etiologies for sporadic miscarriage, 1345-1349
in *LDL receptor* gene as cause of hypercholesterolemia in Taiwan, 1082-1086
of *SLC25A13*, variant clinical courses of 2 patients with neonatal intrahepatic cholestasis with, 1615-1619
- Myoblasts, human, effect of continually high insulin levels on Akt phosphorylation and glucose transport in, 1687-1693
- Myocardial infarction (MI), risk of, association with *APOA1* variant (Val153 greater than Met), 1632-1635
- Myocardial proteins, synthesis of, effect of long-term alcohol administration on, 212-219
- Myocardial substrate selection, influence of calcium-induced workload transitions and fatty acid supply on, 410-420
- Myocardial workload, calcium-induced, and fatty acid supply, influence on myocardial substrate selection, 410-420
- Myocytes, cardiac, effect of elevated glucose on protein synthesis in, 1453-1460
- Myofibrillar proteins, myocardial synthesis of, effect of long-term alcohol administration on, 212-219
- NADPH, *see* Nicotinamide adenine dinucleotide phosphate
- Native Americans, apo C-III bound to apo B-containing lipoproteins and insulin resistance in Cherokee Indian youth, 180-187
- Neck pain, as somatic marker of depression, S1:41-42
- NEFAs (nonesterified fatty acids), *see* Free fatty acids
- Neonatal intrahepatic cholestasis, 2 patients with novel mutation of *SLC25A13*, variant clinical courses, 1615-1619
- Nephrectomy, effects of chronic chromium treatment in uninephrectomized rats, 1243-1249
- Neurobiology, of depression, monoamines, glucocorticoids and aminergic receptors in, S1:25-26
- Neuropeptide Y/agouti-related protein, role in neurophysiologic modulation of obesity, 1207-1208
- Neurosurgery, and depression, S1:28-32
- Neurotransmission, locus for gene-based susceptibility to MDD, S1:17
- Neutrophils
angiotensin II-stimulated oxidative burst in, effect of fluvastatin in hypercholesterolemic patients, 1147-1154

Neutrophils (continued)

- human, TALase-G6PDase supramolecular complex undergoes retrograde trafficking during pregnancy in, 1027-1033
 - in vitro oxidative burst, effects of propylthiouracil and thiamazole, 300-305
- Nicotinamide adenine dinucleotide phosphate (NADPH) oxidase, role in ROS inhibition by MET, 829-834
- Nicotinic receptors, functional, in pancreatic β cells, 247-254
- NIDDM, *see* Non-insulin-dependent diabetes mellitus
- Nitisinone, effect on urinary HGA levels in patients with alkaptonuria, 719-728
- Nitric oxide metabolites (NOx), *see* Plasma nitric oxide metabolites
- Nitric oxide (NO), role of, and intrarenal blood flow distribution in diabetic rats, 788-792
- Nitric oxide (NO) production, effects of red wine consumption on, in NIDDM, 306-313
- Nitric oxide (NO) synthesis, chronic blockade of, resulting in hypoadiponectionemia, 482-487
- L*- ω -Nitro-*L*-arginine methyl ester (L-NAME), effect on regional RBF in diabetic rats, 788-792
- N,N*-Dimethylglycine, and Hcy, and glycine betaine, in patients attending a lipid clinic, 1-14
- NO, *see* Nitric oxide
- NO-1886 (ibrolipim), effect on expression of UCP3 in skeletal muscle and fat accumulation on HF diet-induced obesity, 1587-1592
- Nondiabetic subjects
- healthy, effect of pravastatin on S_1 and adipocytokine levels in, 947-951
 - with MS, targeting CV risk in, 1065-1074
 - overweight, influence of nutrients on cardiac autonomic function, 1290-1296
 - relation of hypercholesterolemia and MS to adhesion of monocytes to cultured endothelium in, 1467-1471
 - Zucker rats, age and muscle-type modulated role of intramyocellular lipids in insulin resistance progression, 522-528
- Nonesterified fatty acids (free fatty acids; NEFAs)
- effects on plasma resistin and insulin resistance in awake rats, 1142-1146
 - elevated concentrations of, effect on human SM gene expression, 952-959
 - and fasting serum TGs and cholesterol, effects of camostat, 623
 - and glucose metabolism, insulin sensitivity indices of, in relation to serum lipids, in obese children and adolescents, 397-402
 - and lactate, glycerol, triglycerides and hormones after 2-day VLCD, 709
 - prolonged exposure, susceptibility of young, LBW men to diabetogenic effects of, 1398-1406
 - see also* Fasting serum free fatty acids
- Non-insulin-dependent diabetes mellitus (NIDDM; type 2 diabetes mellitus)
- altered response of IGFBP-1 to nutritional deprivation in, 275-280
 - apparent and non-apparent atherosclerosis in, LDL size and subclasses as markers of, 227-234
 - association of albuminuria and hypertension with circulating APAAs in, 188-193
 - association of high-sensitivity C-reactive protein with cardiovascular autonomic dysfunction and insulin resistance in, 552-558
 - association between high serum pentosidine concentrations and arterial stiffness and thickness in, 345-350
 - association between K469E polymorphism of *ICAM-1* gene and plasma fibrinogen in, 381-386
 - association of plasma Hcy with serum IL-6 and CP levels in, 809-814
 - with CAD after coronary angioplasty, effect of rosiglitazone on clinical inflammatory responses in, 590-597
 - complicated by hyperlipidemia, effect of low-dose atorvastatin on circulating MCP-1, 1225-1229
 - correlation between adiponectin-leptin ratio and parameters of insulin resistance in, 281-286
 - and CV disease, development from obesity and MS, sex difference in subjects with normal glucose metabolism, 800-808
 - effect of single versus multiple bouts of exercise in women with, 989-994
 - effects of red wine consumption on insulin resistance and endothelial function in, 306-313
 - effects of rosiglitazone and metformin on plasma resistin levels in, 314-320
 - effects of treatment with TZDs on liver function parameters in, 529-532
 - hyperglycemia prevention and gene expression in insulin-sensitive tissues determined by high-density oligonucleotide microarray analysis, effects of exercise training on, 1571-1581
 - and hypertension, effect of rosiglitazone on novel atherosclerotic risk factors in, 1236-1242
 - and impaired glucose tolerance, impact of glucagon response on postprandial hyperglycemia in men with, 1168-1173
 - in Italian subjects, *see* Italian subjects, with NIDDM
 - in Japanese subjects, *see* Japanese NIDDM subjects
 - and MS, preventive effects of soy-based diet supplemented with stevioside in development of, 1181-1188
 - with obesity, *see* Obesity, NIDDM with
 - PM women with, androgen excess as contributor to CV risk profile, 1626-1631
 - in postmenopausal women, *see* Postmenopausal women, with NIDDM

- Non-insulin-dependent diabetes mellitus (NIDDM; type 2 diabetes mellitus) (continued)
 role of insulinemia or sulfonylurea usage in weight gain in patients with, 28-29
- Non-insulin-dependent diabetes mellitus (NIDDM; type 2 diabetes mellitus) microangiopathy, association with serum levels of AGEs and, 1002-1007
- Non-insulin-dependent diabetes mellitus (NIDDM; type 2 diabetes mellitus) nephropathy
 regional RBF distribution and role of NO in, 788-792
 and retinopathy, association with serum levels of AGEs, 1002-1007
- Non-insulin-dependent diabetes mellitus (NIDDM; type 2 diabetes mellitus) retinopathy, and nephropathy, association with serum levels of AGEs, 1002-1007
- Non-insulin-dependent diabetes mellitus (NIDDM; type 2 diabetes mellitus) with obesity
 effect of 2-day VLED on SM insulin sensitivity in patients on insulin therapy, 1669-1678
- Non-obese (lean) subjects
 eNOS content in AT of African American and white American women, 1368-1373
 Japanese, effects of thorough mastication on postprandial plasma glucose concentrations, 1593-1599
 Japanese NIDDM patients, three measures of TNF- α and insulin resistance in nonobese Japanese NIDDM patients, 1297-1301
 and obese (fa/fa) Zucker rats, physiologic difference concerning adiponectin, 995-1001
 versus obese women, ketone body metabolism in, 1542-1545
- Nonthyroidal illness syndrome (NTIS)
 and malnutrition after stroke, 699-704
 and PCM, effect on disability improvement after stroke, 702
- Normal glucose metabolism, development of NIDDM and CV disease from obesity and MS in subjects with, 800-808
- Normal triglycerides (TGs), and IR, relation to PH-LPL activity in African Americans, 902-909
- NOx, *see* Plasma nitric oxide metabolites
- NTIS, *see* Nonthyroidal illness syndrome
- Nutrient intake ratios, and serum Lp-lipid profile and CHO intake, effects of beef versus soy-based food supplements in older men who perform RT, 769-774
- Nutrients, influence on cardiac autonomic function in nondiabetic overweight subjects, 1290-1296
- Nutritional deprivation, in NIDDM, altered response of insulin-like growth factor-binding protein 1 to, 275-280
- Nutritional status, effect of dysphagia on, 701
- OA, *see* Oleic acid
- Obesity
 in African Americans, effect of modest weight loss on insulin action, 960-965
 children, high-sensitive CRP, TNF- α and cardiovascular risk factors before and after weight loss in, 1155-1161
- Obesity (continued)
 comparison of immediate response to meals enriched with *cis*- or *trans*-FAs in individuals with differing FABP2 genotype, 1652-1658
 current knowledge in neurophysiologic modulation of, 1202-1217
 dietary, phenotype-based treatment of, differential effects of fenofibrate, 421-429
 effect on TGF β 1 release by human AT, 1546-1551
 effects of aerobic exercise training on insulin sensitivity, body weight, body fat, adiponectin and inflammatory markers, 1472-1479
 eNOS content in AT of African American and white American women, 1368-1373
 HF diet-induced, effect of NO-1886 on expression of UCP3 in skeletal muscle and fat accumulation, 1587-1592
 and hypertension and IDDM, levels of sICAM-1, sVCAM-1 and selectins in children and adolescents with, 1020-1026
 insulin sensitivity indices of glucose and free fatty acid metabolism, in relation to serum lipids in, 397-402
 versus lean women, ketone body metabolism in, 1542-1545
 and MS, development into NIDDM and CV disease, sex difference in subjects with normal glucose metabolism, 800-808
 nondiabetic subjects with, influence of nutrients on cardiac autonomic function, 1290-1296
 obese (fa/fa) and lean Zucker rats, physiologic difference concerning adiponectin, 995-1001
 prepubertal
 and pubertal youth, altered spontaneous secretion and metabolic clearance of GH in, 1374-1383
 Turkish children, endothelial activation and inflammation in, 1384-1389
 recent onset of, effect of high-carbohydrate meal on postprandial thermogenesis and sympathetic nervous system activity in boys with, 430-438
 severe, impaired response of cardiac ANS to glucose load in, 966-974
 skeletal muscle cell IR in, analysis of insulin receptor activation and glucose transport, 598-603
- Obesity, NIDDM with
 effect of VLCD and withdrawal of blood glucose-lowering therapies on endogenous glucose production in, 705-712
 effects of synthetic PI camostat on dyslipidemia, insulin secretory defects and histological abnormalities of pancreas in, 619-627
- Offspring, body composition and metabolism of, effect of high-fat fed maternal rats, 500-507
- OGTT, *see* Oral glucose tolerance test
- Oil, containing MCTs, effects of dietary DAG on, in patient with LPL deletion, 67-71

- OKG, *see* Ornithine α -ketoglutarate
- Older (elderly) subjects
- suicide in, S1:43
 - underdiagnosis and undertreatment of subsyndromal depression in, S1:39–44
 - versus younger men with similar and smaller BMI and waist circumferences, oxidative and nonoxidative glucose disposal in, 748–755
- Oleic acid (OA)
- effects on vascular reactivity in healthy adult rats, 1600–1609
 - exogenous, effect on intestinal assembly and secretion of apo B48-containing Lp particles in fasting state, 695
- OLETEF, *see* Otsuka Long-Evans Tokushima Fatty
- Omega (ω)-3 fatty acids (FAs), systematic review and meta-analysis of impact on selected arrhythmia outcomes in animal models, 1557–1565
- Omental adipocytes, versus SC adipocytes, basal and insulin-stimulated glucose transport, 781–785
- OPG, *see* Osteoprotegerin
- Opossums, exhibiting high and low lipemic responses to dietary cholesterol and fat, LDL apo B production in, 1075–1081
- Oral glucose tolerance test (OGTT)
- contribution in prevalence of subjects with metabolic syndrome, 542–547
 - endurance training-induced changes in insulin response to, association with PPAR γ 2 Pro12Ala genotype, in men and women, 97–102
 - of male offspring of high-fat fed rats, 500–507
- Oral synthetic estrogen, versus transdermal natural estrogen, effect on Lp(a), Hcy, and CRP levels in PM women, 72–78
- Orexin, role in neurophysiologic modulation of obesity, 1208
- Orlistat, in targeting CV risk in nondiabetic patients with MS, 1065–1074
- Ornithine α -ketoglutarate (OKG), metabolism in healthy rat in postabsorptive state, 1108–1114
- Osteoprotegerin (OPG)
- circulating, *see* Circulating osteoprotegerin
 - plasma, *see* Plasma osteoprotegerin
- Otsuka Long-Evans Tokushima Fatty (OLETF), rats, effects of TZDs on regulations of AQP and GlyK in, 1282–1289
- Overweight, *see* Obesity
- Oxidase, NADPH, role in ROS inhibition by MET, 829–834
- Oxidation, and uptake of FA, effect of short-term leptin treatment in muscle of high fat-fed rats, 1218–1224
- Oxidation, CHO
- of combined glucose and sucrose ingestion during exercise, 610–618
 - during human walking, 793–799
- Oxidative burst
- angiotensin II-induced, sensitivity to fluvastatin in neutrophils of hypercholesterolemic patients, 1147–1154
 - of neutrophil, in vitro, effects of propylthiouracil and thiamazole, 300–305
- Oxidative stress
- in Japanese Americans, protective effect of adiponectin against, 194–199
 - role in liver adaptations to sucrose and fructose, 1195
- Oxidized low density-lipoprotein (OxLDL), and plasma antioxidant capacity in men after short-term cranberry juice consumption, 856–861
- OxLDL, *see* Oxidized low density-lipoprotein
- PAI, *see* Plasminogen activator inhibitor
- PAI-1, *see* Plasminogen activator inhibitor-1
- Palmitate
- chain length, spatial configuration and degree of unsaturation, dose-dependent effects on glucagon secretion, 1329–1336
 - metabolism, effect of short-term leptin treatment on FA uptake and oxidation in muscle of high fat-fed rats, 1222
- PAMBA, *see* *p*-aminomethylbenzoic acid
- p*-Aminomethylbenzoic acid (PAMBA), effect of aromatic ring-containing drugs on carnitine biosynthesis, 1582–1586
- Pancreas, histological abnormalities of, and dyslipidemia and insulin secretory defects, effects of synthetic PI camostat in genetically obese and diabetic rats, 619–627
- Pancreatic beta cells, *see* Beta (β) cells
- Pancreatic insulin content, and serum insulin, glucose response to IVGTT and IR, effects of camostat, 623
- Paraoxonase 1 (PON1) gene, human serum, effect of pitavastatin on transactivation of, 142–150
- Parasympathetic dysfunction (PSD), cardiac, relation to nutrient intake in nondiabetic overweight subjects, 1290–1296
- P12A variant, in PPAR γ , association with insulin resistance in Hispanic and non-Hispanic white women, 1552–1556
- PCM, *see* Protein-energy malnutrition
- PCOS, *see* Polycystic ovary syndrome
- PCs, *see* Phosphatidylcholines
- PKD4, *see* Pyruvate dehydrogenase kinase 4
- Pedigree, human, genetic regulation of variation of circulating IGF-I, IGF-BPs and leptin in, 975–981
- Pentosidine, high serum concentrations, association with arterial stiffness and thickness, in NIDDM, 345–350
- Peptide YY, role in neurophysiologic modulation of obesity, 1210
- Percutaneous coronary intervention, NIDDM subjects with CAD after, effect of rosiglitazone on clinical inflammatory responses in, 590–597

- Peripheral nervous system, role in neurophysiologic modulation of obesity, 1209
- Peroxisome proliferator-activated receptor(s) (PPARs), expression in obese (fa/fa) and lean Zucker rats, 999
- Peroxisome proliferator-activated receptor γ (PPAR γ)
- C161→T polymorphism, association with insulin resistance in Hispanic and non-Hispanic white women, 1552-1556
 - mutant, effect of transgenic expression in liver on fasting-induced and HF diet-induced steatosis in mice, 1490-1498
- Peroxisome proliferator-activated receptor γ (PPAR γ) agonist
- FK-614, effects on peripheral glucose utilization and hepatic insulin extraction in alloxan-induced diabetic dogs, 1250-1258
 - rosiglitazone, effect on clinical inflammatory responses in NIDDM with CAD after coronary angioplasty, 590-597
- Peroxisome proliferator-activated receptor γ 2 (PPAR γ 2)
- gene, Pro12Ala polymorphism of, inverse effects on measures of adiposity over 15 years in African Americans and whites, 910-917
- Peroxisome proliferator-activated receptor- γ 2 (PPAR γ 2)
- Pro12Ala genotype, association with endurance training-induced changes in insulin response to oral glucose, in men and women, 97-102
- Pharmacodynamic analysis, and pharmacokinetics of lanreotide Autogel, 1276-1281
- Pharmacokinetics, and population pharmacodynamic analysis of lanreotide Autogel, 1276-1281
- Phenotype-based treatment, of dietary obesity, differential effects of fenofibrate, 421-429
- Phenylketonuria (PKU), effect of diet on plasma adiponectin concentrations, endothelial dysfunction and atherogenesis in patients with, 1350-1355
- Phosphatidylcholines (PCs), in free form, cholesterol efflux in the presence of, 1091
- Phosphatidylinositol 3-kinase (PI3K)
- activation of, and protein synthesis in cultured cardiac myocytes, effect of elevated glucose, 1453-1460
 - effect of continually high insulin levels on Akt phosphorylation and glucose transport in human myoblasts, 1687-1693
 - signaling pathway, inhibition leads to decreased insulin-stimulated adiponectin secretion from 3T3-L1 adipocytes, 1636-1643
- Physical stress, serum cortisol responses to, effect of corticotropin supplementation, 657-668
- Physiologic difference, between obese (fa/fa) and lean Zucker rats concerning adiponectin, 995-1001
- Phytosterols (PS), and very long chain fatty acids (policosanols), effects on plasma lipid levels and cholesterol biosynthesis, 508-814
- PI3K, *see* Phosphatidylinositol 3-kinase
- Pioglitazone
- effect on body composition and EE, 24-32
 - effect on plasma adiponectin, in chronic blockade of NO synthesis, 482-487
 - for hyperglycemia after androgen-deprivation therapy for prostate cancer, 55-59
- PIs, *see* Protease inhibitors
- Pitavastatin, effect on transactivation of human serum *PON1* gene, 142-150
- Pituitary function, in male and female hypoleptinemic lipodystrophic patients, long-term effect of recombinant methionyl human leptin therapy on, 255-263
- Pituitary gigantism, cardiac and metabolic effects of chronic GH and IGF-1 excess in young adults with, 1174-1180
- PKB/Akt, *see* Protein kinase B
- PKU, *see* Phenylketonuria
- Plant sterols, absorption and plasma concentrations of, in HC, effect of SCH 58235 and high plant sterol-enriched diet on, 38-48
- Plasma adipocytokines, relationship to body fat, in end-stage renal disease, 330-334
- Plasma adiponectin
- in chronic blockade of NO synthesis, 482-487
 - and glucose production, in patients infected with *Plasmodium falciparum*, 60-66
 - increased in poorly controlled patients with phenylketonuria normalize with strict diet, 1350-1355
 - physiologic difference between obese (fa/fa) and lean Zucker rats concerning, 995-1001
 - reciprocal association with insulin resistance and C-reactive protein, in clinically healthy men, 439-444
 - relationship to body fat, in end-stage renal disease, 330-334
- Plasma alanine aminotransferase (ALT), effects of ASO targeted against GCCR, 853
- Plasma alkaline phosphatase (ALP), effects of ASO targeted against GCCR, 853
- Plasma antioxidant capacity, and OxLDL levels in men after short-term cranberry juice consumption, 856-861
- Plasma apolipoprotein C-III (apo C-III), variations in levels of, correlation to TG response to high MUFA diet and high-CHO diet, 1390-1397
- Plasma aspartate aminotransferase (AST), effects of ASO targeted against GCCR, 853
- Plasma carnitine, in patients attending a lipid clinic, 1-14
- Plasma catecholamine, and blood pressure changes during prolonged hyperinsulinemia, 391-396
- Plasma cholesterol (C)
- effects of soy-rich diet and stevioside in NIDDM rats, 1186
- Plasma corticosteroid-binding globulin (CBG), effect on circulating adiponectin-cortisol relationship, in men and women, 584-589

- Plasma corticosterone, effects of ASO targeted against GCCR, 853
- Plasma corticotropin, effects of ASO targeted against GCCR, 853
- Plasma cortisol, and leptin levels, during HC induction of hyperleptinemia, 878
- Plasma C-reactive protein (CRP), in NIDDM with CAD after coronary angioplasty, effect of rosiglitazone, 590-597
- Plasma dehydroepiandrosterone sulfate (DHEA-S), declining levels, and age, contribution to features of MS in men, 1034-1040
- Plasma epinephrine, at rest and during exercise in middle-aged and young athletes versus sedentary men, 1416
- Plasma estradiol, insulin and glucose, in obese and lean African American and white American women, 1370-1371
- Plasma fibrinogen
association with K469E polymorphism of *ICAM-1* gene, in NIDDM, 381-386
and other coronary risk factors, 165-170
in patients with NIDDM complicated with hyperlipidemia, effect of low-dose atorvastatin, 1225-1229
- Plasma ghrelin
effects of FFAs during euglycemic-hyperinsulinemic clamping, 1144
fasting, relationship to metabolic parameters in children and adolescents, 925-929
- Plasma glucagon, effects of ASO targeted against GCCR, 853
- Plasma glucose
estradiol and insulin in obese and lean African American and white American women, 1370-1371
fasting, *see* Fasting plasma glucose
immediate and long-term effects of intranasal insulin administration, 1359
and insulin levels in HC induction of hyperleptinemia, 877-878
levels before and after exercise in insulin-treated STZ-DM fasted rats, 1424
postprandial, *see* Postprandial plasma glucose
at rest and during exercise in middle-aged and young athletes versus sedentary men, 1416
- Plasma glycine betaine, in patients attending a lipid clinic, 1-14
- Plasma high-density lipoprotein-cholesterol (HDL-C), sex-specific effect of *APOA1* variant (Val153 > Met), 1632-1635
- Plasma homocysteine (Hcy)
association with serum IL-6 and CP levels in NIDDM patients, 809-814
concentrations in ESRD, effects of hydroxycobalamin versus cyanocobalamin, 1362-1367
in patients attending a lipid clinic, 1-14
- Plasma homogentisic acid (HGA), in alkaptonuria patients, effect of nitisinone on, 722
- Plasma hormones, and substrates at rest and during exercise in middle-aged and young athletes versus sedentary men, 1416
- Plasma 3-hydroxy-butylate (3-HB), effects of HX531 in KK-A^y mice, 574-576
- Plasma hypoxanthine (Hx), influence of allopurinol on urinary purine loss after repeated sprint exercise in man, 1271-1272
- Plasma inosine, influence of allopurinol on urinary purine loss after repeated sprint exercise in man, 1271-1272
- Plasma insulin
changes in concentration, cardiac ANS response to glucose load in severe obesity, 966-974
effect of modest lifestyle intervention in obese African Americans, 960-965
in elderly versus younger men with similar and smaller BMI and waist circumferences, contributions of oxidative and nonoxidative glucose disposal, 748-755
and estradiol and glucose, in obese and lean African American and white American women, 1370-1371
levels before and after exercise in insulin-treated STZ-DM fasted rats, 1424
at rest and during exercise in middle-aged and young athletes versus sedentary men, 1416
- Plasma insulin immunoreactivity, effect of vagal nerve stimulation, 581
- Plasma lactic acid
levels before and after exercise in insulin-treated STZ-DM fasted rats, 1424
at rest and during exercise in middle-aged and young athletes versus sedentary men, 1416
- Plasma leptin
in children with short stature, effect of insulin administration, 862-865
and leptin resistance, effect of HX531 in KK-A^y mice under normal dietary conditions, 573-578
- Plasma lipids, and cholesterol biosynthesis, effects of very long chain fatty acids (policosanols) and phytosterols, 508-814
- Plasma lipoprotein(a) [Lp(a)], metabolism of apolipoproteins (a) and B-100 within, 361-369
- Plasma low-density lipoprotein-cholesterol (LDL-C), relation to monocyte adhesion to cultured endothelium in nondiabetic subjects, 1467-1471
- Plasma malondialdehyde (MDA), levels, effects of α -lipoic acid, 817-818
- Plasma membrane sialidase (PMS), activities and RSRs in rat liver, effects of chronic ethanol feeding, 1056-1064
- Plasma monocyte chemoattractant protein-1 (MCP-1), in NIDDM with CAD after coronary angioplasty, effect of rosiglitazone, 590-597

- Plasma nitric oxide metabolites (NOx), effect of HF meal in subgroup of young healthy subjects, 1340-1341
- Plasma *N,N*-dimethylglycine, in patients attending a lipid clinic, 1-14
- Plasma osteoprotegerin (OPG), and (s)RANKL, effect of endurance running on, 935-938
- Plasma plasminogen activator inhibitor-1 (PAI-1), levels in patients with NIDDM complicated with hyperlipidemia, effect of low-dose atorvastatin, 1225-1229
- Plasma resistin
effects of rosiglitazone and metformin on, in NIDDM, 314-320
and insulin resistance, effects of FFAs in awake rats, 1142-1146
- Plasma substrates, and hormones at rest and during exercise in middle-aged and young athletes versus sedentary men, 1416
- Plasma testosterone, declining levels, and age, contribution to features of MS in men, 1034-1040
- Plasma total homocysteine (tHcy), maximum reduction in ESRD, dose regimen of cobalamin for, 835-840
- Plasma triglyceride(s) (TGs)
correlation with DNA variants in *apo A5* gene in Chinese, 568-572
effects of soy-rich diet and stevioside in NIDDM rats, 1186
role in pathophysiological subsets of preeclampsia, 1504-1509
- Plasma tyrosine, after nitisinone treatment of alkaptonuria patients, 722-723
- Plasma uric acid, influence of allopurinol on urinary purine loss after repeated sprint exercise in man, 1271-1272
- Plasminogen activator inhibitor (PAI), high activity, and *factor V Leiden* gene mutation and high factor VIII, etiologies for sporadic miscarriage, 1345-1349
- Plasminogen activator inhibitor-1 (PAI-1)
plasma, *see* Plasma plasminogen activator inhibitor-1
release by human adipocytes, effect of insulin on, 220-226
serum, *see* Serum plasminogen activator inhibitor-1
- Plasmodium falciparum* infection, glucose production and plasma adiponectin in, 60-66
- PMS, *see* Plasma membrane sialidase
- PM women, *see* Postmenopausal women
- Policosanols, and phytosterols, effects on plasma lipid levels and cholesterol biosynthesis, 508-814
- Polycystic ovary syndrome (PCOS), and IR, metformin and diet benefits in women with, 113-121
- Polymorphism
Arg211His, of *GPR40* gene, contribution to variation of insulin secretory capacity in Japanese men, 296-299
AvalI, of *LDL-receptor* gene, and baseline LDL-C and Lp(a), influence on LDL-C response to pravastatin, 741-747
- Polymorphism (continued)
C161→T, in PPAR γ , association with insulin resistance in Hispanic and non-Hispanic white women, 1552-1556
functional, of Z-DNA-forming motif of promoter of *SLC11A1* gene, association with IDDM in Japanese, 628-633
G82S and G1704T, of *RAGE* gene, association with microalbuminuria, in Japanese NIDDM patients, 488-491
K469E, of *ICAM-1* gene, association with plasma fibrinogen level in NIDDM, 381-386
Pro12Ala, of *PPAR γ 2 gene, inverse effects on measures of adiposity over 15 years in African Americans and whites, 910-917
single nucleotide, *see* Single nucleotide polymorphisms
PON1, *see* *Paraoxonase 1**
- Population pharmacodynamic analysis, and pharmacokinetics of lanreotide Autogel, 1276-1281
- Postabsorptive state, OKG metabolism in healthy rat in, 1108-1114
- Postheparin-lipoprotein lipase (PH-LPL), activity of, relation to IR and normal TGs in African Americans, 902-909
- Posthepatic insulin clearance rate, and hepatic insulin extraction, in proportion to insulin resistance, in HIV infection with lipodystrophy, 171-179
- Postmenopausal (PM) women
asthmatic, treated with inhaled corticosteroids, effects of alendronate on BMD and bone metabolic markers in, 85-90
effect of HRT on serum adiponectin levels, 1610-1614
effects of types of HRT on Lp(a), Hcy, and CRP levels in, 72-78
Korean, association among serum ferritin, ALT levels and MS in, 1510-1514
with primary hypercholesterolemia, effects of raloxifene and low-dose simvastatin coadministration on serum lipids, 939-946
- Postmenopausal (PM) women, with NIDDM
androgen excess as contributor to CV risk profile, 1626-1631
effects of diet and/or exercise on adipocytokine and inflammatory cytokine levels in, 866-875
relationship between blood lactate concentration and substrate utilization during exercise in, 1102-1107
- Postprandial apolipoprotein B-48 clearance, effect of LPL S447X variant, 1499-1503
- Postprandial control, of EGP, effects of repaglinide and repaglinide on, 79-84
- Postprandial hyperglycemia
impact of glucagon response in men with impaired glucose tolerance and NIDDM, 1168-1173

- Postprandial hyperglycemia (continued)
and subjective abdominal symptoms, effects of voglibose and acarbose on, 387-390
- Postprandial insulin sensitivity (S_i), modifying effect of vagal nerve stimulation on, 579-583
- Postprandial lipemia (lipidemia), dose-response effect of resistance exercise, 756-763
- Postprandial (PP) glucose, and lipids, comparison of immediate response to meals enriched with *cis*- or *trans*-FAs in overweight individuals with differing FABP2 genotype, 1652-1658
- Postprandial (PP) lipids, and glucose, comparison of immediate response to meals enriched with *cis*- or *trans*-FAs in overweight individuals with differing FABP2 genotype, 1652-1658
- Postprandial (PP) plasma glucose, effects of thorough mastication in nonobese Japanese subjects, 1593-1599
- Postprandial thermogenesis, and sympathetic nervous system activity, in boys with recent onset of obesity, effect of high-carbohydrate meal, 430-438
- Postprandial triglycerides (TGs), and IR in primary renal disease, 821-828
- POWER study, *see* Premenopause, Osteopenia/Osteoporosis, Women, Alendronate, Depression study
- PPAR γ , *see* Peroxisome proliferator-activated receptor γ
- PPAR γ 2, *see* Peroxisome proliferator-activated receptor- γ 2
- PPARs, *see* Peroxisome proliferator-activated receptors
- Pravastatin
and atorvastatin and fenofibrate, comparative study of beneficial effects in types IIa and IIb hyperlipidemia patients, 677-681
effect on S_i and adipocytokine levels in healthy nondiabetic patients, 947-951
LDL-C response to, influence of baseline LDL-C, Lp(a) and *Aval*1 polymorphism of *LDL-receptor* gene, 741-747
- Preeclampsia, atherogenic lipoprotein subfraction profile in preeclamptic women with and without high TGs, 1504-1509
- Pregnancy
depression during, S1:50
and lactation, high saturated fat versus high CHO diet during, effects on insulin sensitivity in offspring rats, 1316-1322
MS at follow-up in women with and without GDM, 1115-1121
TALase-G6PDase supramolecular complex undergoes retrograde trafficking in human neutrophils during, 1027-1033
- Preheparin serum lipoprotein lipase (LPL), correlation with adiponectin-leptin ratio, in NIDDM, 281-286
- Premenopausal women
versus PM women, effect of HRT on serum adiponectin levels, 1610-1614
- Premenopausal women (continued)
younger, with MDD, abdominal AT and serum levels of prothrombic factors in, 918-924
- Premenopause, Osteopenia/Osteoporosis, Women, Alendronate, Depression (POWER) study, 918-924
- Premenstrual syndrome, and premenstrual dysphoric disorder, S1:49-50
- Prepubertal boys, racial difference in circulating SHBG levels in, 91-96
- Prevalence
of chronic infections, in relation to CRP levels, in subjects with hypoalbuminoproteinemia, 33-37
of metabolic syndrome, contribution of impaired glucose tolerance test in, 542-547
of premenstrual syndrome and premenstrual dysphoric disorder, S1:49
- Primary hypercholesterolemia, effects of raloxifene and low-dose simvastatin coadministration on serum lipids in PM women with, 939-946
- Primary renal disease, role of CrCl and body composition in development of IR in, 821-828
- Pro12Ala polymorphism, of *PPAR* γ 2 gene, inverse effects on measures of adiposity over 15 years in African Americans and whites, 910-917
- Prolactin-releasing peptide, role in neurophysiologic modulation of obesity, 1209
- Proliferative retinopathy, association with circulating APAAs in NIDDM, 188-193
- Prolonged hyperinsulinemia, blood pressure changes and plasma catecholamine levels during, 391-396
- Propylthiouracil, and thiamazole, effects on in vitro neutrophil oxidative burst, 300-305
- Prostate cancer, hyperglycemia after androgen-deprivation therapy for, and effect of pioglitazone, 55-59
- Protease inhibitors (PIs)
HAART based on, effect on thigh SAT gene expression, 561-567
synthetic, effects on dyslipidemia, insulin secretory defects and histological abnormalities of pancreas in genetically obese and diabetic rats, 619-627
- Protein, *see* Dietary protein
- Protein anabolism, in response to insulin, influence of sex, 1529-1535
- Protein-energy malnutrition (PCM)
effect on thyroid hormonal parameters after stroke, 701-702
and nonthyroidal illness syndrome after stroke, 699-704
and NTIS, effect on disability improvement after stroke, 702
- Protein kinase B (PKB/Akt), activity, and protein synthesis in cultured cardiac myocytes, effect of elevated glucose, 1453-1460

- Protein supplementation, combined with resistance training, effect on muscle fiber size and muscle strength, 151-156
- Protein synthesis
of myofibrillar and sarcoplasmic proteins in heart, effect of long-term alcohol administration on, 212-219
response to hypercapnia in rats, 841-847
- Prothrombic factors, serum levels, and abdominal AT in younger, premenopausal women with MDD, 918-924
- PS, *see* Phytosterols
- PSD, *see* Parasympathetic dysfunction
- P-selectin, *see* soluble P-selectin
- Psychiatric illness, association with medical illness, S1:53-54
- Psychosis, postpartum, S1:51
- P33T allele, *IPF-1* gene missense mutation in Italian family with NIDDM and GDM, 983-988
- Puerariae radix*, effect on bone loss in castrated male mice, 1536-1541
- Pulmonary vessel endothelium, role in fibrinogen metabolism, 271-274
- Purine loss, urinary, after repeated sprint exercise in man, influence of allopurinol, 1269-1275
- Pyruvate dehydrogenase kinase 4 (PDK4), gene expression in SM, effects of elevated lipid concentrations, 952-959
- Quadriceps muscle, of male offspring of high-fat fed rats, 500-507
- Quantitative Insulin Sensitivity Check Index (QUICKI)
effect of pravastatin on S_i in healthy nondiabetic patients, 947-951
spectrum of insulin sensitivity in Korean population, 1644-1651
- QUICKI, *see* Quantitative Insulin Sensitivity Check Index
- Race and ethnicity
association of low levels of adiponectin with metabolic syndrome, 476-481
effect of modest weight loss on insulin action in obese African Americans, 960-965
effect of westernized lifestyle on the association between fasting serum nonesterified fatty acids and insulin secretion in Japanese American men, 713-718
eNOS content in AT of obese and lean African Americans and white American women, 1368-1373
inverse effects of *PPAR* γ 2 Pro12Ala polymorphism on longitudinal measures of adiposity in African Americans and whites, 910-917
protective effect of adiponectin against oxidative stress in Japanese Americans, 194-199
in relation to circulating SHBG levels in prepubertal boys, 91-96
see also Hispanic subjects; Italian subjects; Japanese subjects; Korean subjects; Turkish subjects
- Race and ethnicity, insulin resistance and
and apo C-III bound to apo B-containing lipoproteins in Cherokee Indian youth, 180-187
association with C161T polymorphism in *PPAR* γ in Hispanic and non-Hispanic white women, 1552-1556
relationship to PH-LPL activity and normal TGs in African Americans, 902-909
- RAGE* gene, G1704T and G82S polymorphisms of, association with microalbuminuria, in Japanese NIDDM patients, 488-491
- Raloxifene, and low-dose simvastatin, effects of coadministration on serum lipids in PM women with primary hypercholesterolemia, 939-946
- Rapamycin, p38 inhibitor, effect on glucose uptake in insulin-resistant 3T3-L1 adipocytes, 895-901
- Raw vegetable food (RVF), containing high cyclo (his-pro), effect on insulin sensitivity and body weight control, 1480-1489
- RBF, *see* Renal blood flow
- Reactive oxygen species (ROS), intracellular production in aortic ECs, effect of metformin, 829-834
- Recombinant methionyl human leptin therapy, long-term effect on hyperandrogenism, menstrual function, and pituitary function in hypoleptinemic lipodystrophic patients, 255-263
- Recovery, from exercise, substrate availability and transcriptional regulation of metabolic genes in human SM during, 1048-1055
- Rectal temperature, effects of HX531 in KK-A γ mice, 574-576
- Recurrent pregnancy loss (RPL), and miscarriages, thrombophilia and hypofibrinolysis in women with, 1346-1347
- Red wine consumption, effects on insulin resistance and endothelial function in NIDDM, 306-313
- Reflex increases, in adrenal or lumbar SNA induced by hypoglycemia, effect of captopril, 1679-1686
- Relative synthetic rates (RSRs), and activities of various sialidases in rat liver, effects of chronic ethanol feeding, 1056-1064
- Renal blood flow (RBF), regional, and role of NO in diabetic rats, 788-792
- Renal disease, role of CrCl and body composition in development of IR in, 821-828
- Repaglinide, and gliclazide, effects on postprandial control of EGP, 79-84
- Reproductive mood disorders, in women, S1:49-51
- Resistance training (RT)
combined with timed ingestion of protein, effect on muscle fiber size and muscle strength, 151-156
dose-response effect on postprandial lipemia, 756-763
older men who perform, effects of beef versus soy-based food supplements on serum Lp-lipid profile, CHO intake and nutrient intake ratios in, 769-774

- Resistin
 plasma, *see* Plasma resistin
 role in neurophysiologic modulation of obesity, 1206
 serum, *see* Serum resistin
- Respiratory mitochondrial chain, role in ROS inhibition by MET, 829-834
- Retinoid X receptor antagonist, HX531, effect on leptin resistance without increasing plasma leptin level, in KK-A^y mice under normal dietary conditions, 573-578
- Retinopathy
 proliferative, association with circulating APAAs in NIDDM, 188-193
 see also Non-insulin dependent diabetes mellitus retinopathy
- Retrograde trafficking, of TALase-G6PDase supramolecular complex, in human neutrophils during pregnancy, 1027-1033
- Right ventricle (RV) hypertrophy, and hypoxia-induced anorexia, role in lactate transport and monocarboxylate transporter expression in rat muscle, 634-644
- Risk
 for CAD, association with circulating sCD44, 139-141
 see also Risk, for CV disease
- Risk, for CV disease
 association with abdominal AT and serum levels of prothrombic factors in younger, premenopausal women with MDD, 918-924
 and high-sensitive CRP and TNF- α before and after weight loss in obese children, 1155-1161
 of MS, relation to serum ferritin and ALT levels in Korean postmenopausal women, 1510-1514
 relation to serum adiponectin levels, 1610-1614
 role of androgen excess in postmenopausal women with NIDDM, 1626-1631
 targeting in nondiabetic patients with MS, 1065-1074
- Risk factors, for CAD, and fibrinogen plasma concentration, 165-170
- ROS, *see* Reactive oxygen species
- Rosiglitazone (RSG)
 effect on clinical inflammatory responses in NIDDM with CAD after coronary angioplasty, 590-597
 effect on novel atherosclerotic risk factors in patients with NIDDM and hypertension, 1236-1242
 effect on serum Hcy levels, smooth muscle proliferation and intimal hyperplasia, in high methionine-fed Sprague-Dawley rats, 645-652
 effects on regulations of AQP and GlyK in OLETF rats, 1282-1289
 and metformin, effects on plasma concentrations of resistin, in NIDDM, 314-320
- RPL, *see* Recurrent pregnancy loss
- R155P mutation, in *ALADIN* gene, and Allgrove syndrome, diagnosis of adrenal insufficiency in patient with, 200-205
- RSG, *see* Rosiglitazone
- RT, *see* Resistance training
- Running, endurance, effects on plasma OPG and (s)RANKL, 935-938
- RVF, *see* Raw vegetable food
- SA, *see* Stearic acid
- Sarcoplasmic proteins, myocardial synthesis of, effect of long-term alcohol administration on, 212-219
- SAT, *see* Subcutaneous adipose tissue
- Satiety, and hunger, effect of pioglitazone on, 24-32
- Saturated fatty acid(s) (SFA), versus dietary *trans*-FA, effects on adipocyte plasma membrane fatty acid composition and insulin sensitivity, 240-246
- sCD44, *see* Soluble CD44
- SCH 58235, and high-plant-sterol diet, effects on plant sterol absorption and plasma concentrations in hypercholesterolemic WKY rats, 38-48
- Sedentary subjects, versus middle-aged men and young athletes, effects of exercise at moderate and hard intensity on substrate oxidation, 1411-1419
- Selectins, *see specific selectins*
- Serotonin, role in premenstrual syndrome and premenstrual dysphoric disorder, S1:50
- Serotonin-uptake gene, and stress and MDD, S1:17-18
- Sertoli cells, cultured, metabolism of amino acids by, 515-521
- Serum adiponectin
 association with arteriosclerosis obliterans and IHD, 653-656
 effect of postmenopause and HRT on, 1610-1614
 effect of thyroid hormone in Basedow disease, 1461-1466
 inverse correlation with HLC, in Japanese NIDDM subjects, 775-780
- Serum advanced glycation end products (AGEs), association with NIDDM retinopathy and nephropathy, 1002-1007
- Serum cortisol
 relation to adiponectin, effect of CBG in men and women, 584-589
 responses to physical stress, effect of cortisol supplementation, 657-668
- Serum dehydroepiandrosterone sulfate (DHEAS), association with NIDDM progression in Japanese men, 669-676
- Serum factor VIII (fVIII), and abdominal AT in younger premenopausal women with MDD, 918-924
- Serum ferritin, relation to ALT levels and MS in Korean postmenopausal women, 1510-1514
- Serum free thyroxine, lower levels, association with MS in Chinese population, 1524-1528
- Serum glucose, relation to ACP₁ genotype in healthy adult subjects, 891-894

- Serum hepatic enzyme, and waist circumference, and birth weight, of 18-year old sons, associations with their father's and mother's BMI, 466-470
- Serum high-density lipoprotein-cholesterol (HDL-C), effect of HF meal in subgroup of young healthy subjects, 1340
- Serum homocysteine (Hcy), and smooth muscle proliferation and intimal hyperplasia, effect of rosiglitazone in high methionine-fed Sprague-Dawley rats, 645-652
- Serum insulin
effect of HF meal in subgroup of young healthy subjects, 1340
and glucose response to IVGTT, insulin resistance, and pancreatic insulin content, effects of camostat, 623
- Serum interleukin-6 (IL-6), and CP levels, association with plasma Hcy in NIDDM patients, 809-814
- Serum leptin, TNF- α -associated expression of, in dimethylnitrosamine and bile duct ligation-induced liver cirrhosis, 445-452
- Serum lipid profile
relationship to soluble E-selectin, leptin, and insulin resistance, in nonobese Japanese with NIDDM, 376-380
sTNF-R, BMI, leptin and adiponectin in nonobese Japanese NIDDM patients, 1297-1301
- Serum lipids
and colonic microflora, effect of antibiotics on, 103-112
effects of raloxifene and low-dose simvastatin coadministration in PM women with primary hypercholesterolemia, 939-946
physiological relevance of apoE recycling in primary mouse hepatocytes, 1311
in relation to insulin sensitivity indices of glucose and free fatty acid metabolism, in obese children and adolescents, 397-402
- Serum lipoprotein lipase (LPL), preheparin, correlation with adiponectin-leptin ratio, in NIDDM, 281-286
- Serum low-density lipoprotein-cholesterol (LDL-C), effect of HF meal in subgroup of young healthy subjects, 1340
- Serum *para*oxonase 1 (PON1) gene, effect of pitavastatin on transactivation of, 142-150
- Serum pentosidine, association with arterial stiffness and thickness, in NIDDM, 345-350
- Serum plasminogen activator inhibitor-1 (PAI-1), and abdominal AT in younger premenopausal women with MDD, 918-924
- Serum prothrombic factors, and abdominal AT in younger, premenopausal women with MDD, 918-924
- Serum resistin, relationship to anthropometric and metabolic parameters, among healthy subjects, 471-475
- Serum total cholesterol (TC)
association with circulating OPG in healthy Korean women, 49-54
effect of HF meal in subgroup of young healthy subjects, 1340
- Serum triacylglycerols (TAGs), effects of DAG on, in patient homozygous for complete lipoprotein lipase deletion, 67-71
- Serum triglyceride(s) (TGs), effect of HF meal in subgroup of young healthy subjects, 1340
- Serum uric acid
genome-wide search for genes affecting, 1435-1441
high levels, walking time for, and alcohol drinking, possible interactions with eNOS genotype among Japanese, 1302-1308
- sE-selectin, *see* Soluble E-selectin
- Sex
associations with cardiovascular risk factors, and subclinical atherosclerosis, 139-141
and β_2 adrenergic receptor haplotype, effect on resting and terbutaline-stimulated whole body lipolysis, 492-499
differences, effect of CBG on circulating adiponectin-cortisol relationship, 584-589
differences in development of NIDDM and CV disease from obesity and MS, in subjects with normal glucose metabolism, 800-808
differences in plasma Hcy and plasma and urine glycine betaine prediction, in patients attending a lipid clinic, 7, 9
differences in relationship between fasting plasma ghrelin and metabolic parameters in children and adolescents, 925-929
influence on protein anabolic response to insulin, 1529-1535
long-term effect of recombinant methionyl human leptin therapy on pituitary function in male and female hypoleptinemic lipodystrophic patients, 255-263
in relation to PPAR γ 2 Pro12Ala genotype and insulin action, 97-102
specific effect of *APOA1* variant (Val153 > Met) on plasma HDL-C, 1632-1635
see also Men; Women
- Sex hormone-binding globulin (SHBG)
circulating, in prepubertal boys, in relation to racial difference, 91-96
relation to serum adiponectin levels, 1610-1614
- Sex hormones, changes in balance of, in premenstrual syndrome and premenstrual dysphoric disorder, S1:49-50
- Sex steroids, and/or GH, effect on whole-body protein turnover in healthy aged men and women, 1162-1167
- SFA, *see* Saturated fatty acids
- SHBG, *see* Sex hormone-binding globulin
- Shear rates, whole-blood viscosity at, influence of Lp, 764-768
- Short stature, idiopathic, leptin responses to insulin administration in children with, 862-865
- Short-term leptin treatment, effect on FA uptake and oxidation in muscle of high fat-fed rats, 1218-1224

- S₁, *see* Insulin sensitivity
- Sialidase
activities and RSRs in rat liver, effects of chronic ethanol feeding, 1056-1064
see also specific sialidases
- Gal β 1, 4GlcNAc α 2,6-sialyltransferase messenger RNA (2,6-ST mRNA), down-regulation of, effect of ethanol exposure in human liver cell lines, 729-734
- sICAM-1, *see* Soluble intercellular adhesion molecule-1
- Simvastatin, low-dose, and raloxifene, effects of coadministration on serum lipids in PM women with primary hypercholesterolemia, 939-946
- Single nucleotide polymorphisms (SNPs), of *apo A5* gene, association with CHD in Chinese, 568-572
- Skeletal mass, effect of endurance running on plasma OPG and (s)RANKL, 935-938
- Skeletal muscle (SM)
cells, from obese subjects, analysis of insulin-stimulated insulin receptor activation and glucose transport in, 598-603
gene expression
effect of elevated lipid concentrations, 952-959
and hyperglycemia prevention in Zucker diabetic fatty rats, effects of exercise training on, 1571-1581
insulin sensitivity, effect of 2-day VLED in obese NIDDM patients on insulin therapy, 1669-1678
postexercise glycogen repletion in, effect of impaired glucose uptake in insulin-treated STZ-DM fasted rats, 1420-1427
protein synthesis in, effects of acidosis and hypothermia on, 841-847
during recovery from exercise, substrate availability and transcriptional regulation of metabolic genes in, 1048-1055
UCP3 expression in, and fat accumulation, effect of NO-1886 on HF diet-induced obesity, 1587-1592
- SLC11A1* gene, functional polymorphism of Z-DNA-forming motif of promoter of, association with IDDM in Japanese, 628-633
- SLC25A13* gene mutation, variant clinical courses of 2 patients with neonatal intrahepatic cholestasis with, 1615-1619
- SM, *see* Skeletal muscle
- Smooth muscle proliferation, and serum Hcy and intimal hyperplasia, effect of rosiglitazone in high methionine-fed Sprague-Dawley rats, 645-652
- SNA, *see* Sympathetic nerve activity
- SNPs, *see* Single nucleotide polymorphisms
- Soleus muscle, gene expression in, and hyperglycemia prevention in Zucker diabetic fatty rats, effects of exercise training on, 1571-1581
- Soluble CD44 (sCD44), circulating, associations with sex, cardiovascular risk factors, and subclinical atherosclerosis, 139-141
- Soluble endothelial leukocyte adhesion molecule-1, *see* Soluble E-selectin
- Soluble E-selectin
endothelial activation in prepubertal obese Turkish children, 1384-1389
relationship to leptin, triglycerides, and insulin resistance, in nonobese Japanese with NIDDM, 376-380
and sVCAM-1 and sICAM-1, levels in children and adolescents with obesity, hypertension and IDDM, 1020-1026
- Soluble intercellular adhesion molecule-1 (sICAM-1)
endothelial activation in prepubertal obese Turkish children, 1384-1389
and insulin sensitivity in overweight and obese girls, effects of aerobic exercise training, 1472-1479
and sVCAM-1 and selectins, levels in children and adolescents with obesity, hypertension and IDDM, 1020-1026
- Soluble L-selectin, and sVCAM-1 and sICAM-1, levels in children and adolescents with obesity, hypertension and IDDM, 1020-1026
- Soluble P-selectin, and sVCAM-1 and sICAM-1, levels in children and adolescents with obesity, hypertension and IDDM, 1020-1026
- Soluble receptor activator of nuclear factor κ B ligand [(s)RANKL], plasma, and OPG, effect of endurance running on, 935-938
- Soluble tumor necrosis factor receptor(s) (sTNF-R)
and BMI, leptin, adiponectin and serum lipid profile in nonobese Japanese NIDDM patients, 1297-1301
effects of metformin on, 235-239
- Soluble vascular cell adhesion molecule-1 (sVCAM-1)
endothelial inflammation in prepubertal obese Turkish children, 1384-1389
and insulin sensitivity in overweight and obese girls, effects of aerobic exercise training, 1472-1479
and sICAM-1 and selectins, levels in children and adolescents with obesity, hypertension and IDDM, 1020-1026
- Somatic disorders, association with mental illness, S1:53-54
- Soy-based diet
versus beef intake, effects on serum Lp-lipid profiles, CHO intake and nutrient intake ratios, in older men who perform RT, 769-774
supplemented with stevioside, preventive effects on development of MS and NIDDM in Zucker diabetic fatty rats, 1181-1188
- Sporadic miscarriage, etiologies, role of *factor V Leiden* gene mutation, high factor VIII and high PAI activity, 1345-1349
- Sprint exercise, repeated, influence of allopurinol on urinary purine loss after, 1269-1275
- (s)RANKL, *see* Soluble receptor activator of nuclear factor κ B ligand
- SREBP-1c, *see* Sterol regulatory element binding protein 1c
- SSPG, *see* Steady-state plasma glucose

- Statins, *see* Hydroxymethyl glutaryl coenzyme A reductase inhibitors
- Steady-state plasma glucose (SSPG), relationship between IMGU and excess adiposity as determined by BMI and WC, 1323-1328
- Stearic acid (SA), effects on vascular reactivity in healthy adult rats, 1600-1609
- Steatosis, fasting-induced and HF diet-induced, effect of transgenic expression of mutant PPAR γ in liver, 1490-1498
- Sterol regulatory element binding protein 1c (SREBP-1c), modification of gene expression by leptin, effect on lipogenic enzyme expression in WAT of aging rats, 1041-1047
- Stevioside, soy-based diet supplemented with, preventive effects on development of MS and NIDDM in Zucker diabetic fatty rats, 1181-1188
- sTNF-R, *see* Soluble tumor necrosis factor receptor(s)
- Streptozotocin-induced diabetes mellitus (STZ-DM), insulin-treated fasted rats, effect of impaired glucose uptake on postexercise glycogen repletion in SM, 1420-1427
- Stress
effect on prefrontal cortex and amygdala, S1:21-22
and genes, relation to depressive disorders, S1:16-19
physical, serum cortisol responses to, effect of cortitrol supplementation, 657-668
- Stroke
disability improvement after, effect of PCM and NTIS, 702
malnutrition and nonthyroidal illness syndrome after, 699-704
thyroid hormonal parameters after, effect of PCM on, 701-702
- STZ-DM, *see* Streptozotocin-induced diabetes mellitus
- Subcutaneous adipose tissue (SAT)
accumulation, and UCP3 expression in skeletal muscle, effect of NO-1886 on HF diet-induced obesity, 1587-1592
ANP- and adrenergic receptor-dependent lipolytic pathways in, differential regulation, 122-131
effect of TZDs on, 24-32
thigh, alterations in gene expression of, effect of PI-based HAART, 561-567
- Subcutaneous (SC) insulin, versus buccal spray insulin, in IDDM patients, 930-934
- Subjective abdominal symptoms, and postprandial hyperglycemia, effects of voglibose and acarbose on, 387-390
- Substrate availability, and transcriptional regulation of metabolic genes in human SM during recovery from exercise, 1048-1055
- Substrate oxidation, effects of exercise at moderate and hard intensity in middle-aged and young athletes versus sedentary men, 1411-1419
- Substrate utilization, relationship to blood lactate concentration during exercise in NIDDM postmenopausal women, 1102-1107
- Subsyndromal depression, underdiagnosis and undertreatment in elderly, S1:39-44
- Sucrose
combined ingestion with glucose, oxidation during exercise, 610-618
and fructose, hepatic adaptations to, 1189-1201
- Suicide, in elderly, S1:43
- Sulfonylurea(s) usage, role in weight gain in NIDDM patients, 28-29
- Supplementation
carbohydrate, versus isoenergetic protein ingestion, combined with resistance training, effect on muscle fiber size and muscle strength, 151-156
carnitine, effect on lipoprotein metabolism and LDL atherogenicity, 1135
with cortitrol, effect on serum cortisol responses to physical stress, 657-668
 ω -3 fatty acids, systematic review and meta-analysis of impact on selected arrhythmia outcomes in animal models, 1557-1565
flavonoid-rich cranberry juice, effect on plasma antioxidant capacity and OxLDL levels in men, 856-861
with α -lipoic acid, effects on exercise-induced blood coagulation activation, 815-820
with soy-based food versus beef intake, effects on serum Lp-lipid profiles, CHO intake and nutrient intake ratios, in older men who perform RT, 769-774
with stevioside and soy-based diet, preventive effects on development of MS and NIDDM in Zucker diabetic fatty rats, 1181-1188
- Supramolecular complex, of TALase-G6PDase, retrograde trafficking in human neutrophils during pregnancy, 1027-1033
- sVCAM-1, *see* Soluble vascular cell adhesion molecule-1
- S447X variant, of LPL, effect on postprandial apo B-48 clearance, 1499-1503
- Sympathetic nerve activity (SNA), adrenal or lumbar, hypoglycemia-induced, effect of captopril on reflex increases in, 1679-1686
- Sympathetic nervous system activity, and postprandial thermogenesis, in boys with recent onset of obesity, effect of high-carbohydrate meal, 430-438
- Systolic blood pressure (BP), endothelial activation and inflammation in prepubertal obese Turkish children, 1384-1389
- TAGs, *see* Serum triacylglycerols
- Taiwan, hypercholesterolemia in, caused by mutations in *LDL receptor* gene, 1082-1086
- TALase, *see* Transaldolase
- Terbutaline-stimulated, and resting whole body lipolysis, effect of sex and β_2 adrenergic receptor haplotype on, 492-499
- Testosterone, declining plasma levels, and age, contribution to features of MS in men, 1034-1040

- Testosterone secretion, in adolescent boys with β -thalassemia major versus CDGP, effects of hCG therapy on, 18–20
- Texturized vegetable protein (TVP), versus beef intake, effects on serum Lp-lipid profiles, CHO intake and nutrient intake ratios, in older men who perform RT, 769–774
- TGF- β 1, *see* Tumor growth factor β 1
- TGs, *see* Triglyceride(s)
- β -Thalassemia major, versus CDGP, human chorionic gonadotropin therapy in adolescent boys with, 15–23
- tHcy, *see* Plasma total homocysteine
- Thermogenesis, postprandial, and sympathetic nervous system activity, in boys with recent onset of obesity, effect of high-carbohydrate meal, 430–438
- Thiamazole, and propylthiouracil, effects on *in vitro* neutrophil oxidative burst, 300–305
- Thiazolidinedione(s) (TZDs)
- effect on liver function parameters, in NIDDM patients, 529–532
 - effect on visceral, subcutaneous, and total AT, 24–32
 - effects on regulations of AQP and GlyK in OLETF rats, 1282–1289
 - for hyperglycemia after androgen-deprivation therapy for prostate cancer, 55–59
 - rosiglitazone, effect on novel atherosclerotic risk factors in patients with NIDDM and hypertension, 1236–1242
- Thigh subcutaneous adipose tissue (SAT), alterations in gene expression of, effect of PI-based HAART, 561–567
- THP-1 mononuclear phagocytes, effect of iron loading on cholesterol accumulation and macrophage scavenger receptor I expression in, 453–459
- 3T3-L1 adipocytes
- insulin-resistant, effect of p38 MAP kinase inhibition on glucose uptake in, 895–901
 - insulin-stimulated adiponectin secretion from, effect of inhibition of PI3K signaling pathway, 1636–1643
- Thrombophilia, and hypofibrinolysis in women with miscarriages and RPL, 1346–1347
- Thyroid hormonal parameters, after stroke, effect of PCM on, 701–702
- Thyroid hormone, effect on serum adiponectin levels in Basedow disease, 1461–1466
- Thyroid-stimulating hormone receptor antibody (TRAb), association with serum adiponectin levels in Basedow disease, 1461–1466
- Tissue protein synthesis, effects of acute respiratory acidosis and hypothermia on, 841–847
- Tissue weight, effects of VLCFAs (policosanols) and phytosterols on, 511–512
- TNF- α , *see* Tumor necrosis factor- α
- Total adipose tissue (AT), effect of TZDs on, 24–32
- Total homocysteine (tHcy), *see* Plasma total homocysteine
- TRAb, *see* Thyroid-stimulating hormone receptor antibody
- Transaldolase (TALase), supramolecular complex with G6PDase, retrograde trafficking in human neutrophils during pregnancy, 1027–1033
- Transcription activation, of human serum *PON1* gene, effect of pitavastatin on, 142–150
- Transcriptional regulation, and substrate availability of metabolic genes in human SM during recovery from exercise, 1048–1055
- Transdermal natural estrogen, versus oral synthetic estrogen, effect on Lp(a), Hcy, and CRP levels in PM women, 72–78
- Trans-fatty acids (FA), dietary, effects on adipocyte plasma membrane fatty acid composition and insulin sensitivity, 240–246
- Translocation, of GLUT4, effect of 2-day VLED on SM insulin sensitivity in obese NIDDM patients on insulin therapy, 1669–1678
- Triacylglycerols (TAGs), *see* Serum triacylglycerols
- Triglyceride(s) (TGs)
- accumulation in tissues of diet-induced insulin-resistant hyperlipidemia, association with increased diacylglycerol acyltransferase activity, 403–409
 - correlation with adiponectin-leptin ratio, in NIDDM, 281–286
 - effect of 2-day VLED on SM insulin sensitivity in obese NIDDM patients on insulin therapy, 1669–1678
 - muscle, effect of short-term leptin treatment on FA uptake and oxidation in high fat-fed rats, 1218–1224
 - and nonesterified fatty acids, lactate, glycerol and hormones after 2-day VLCD, 709
 - normal, *see* Normal triglycerides
 - plasma, *see* Plasma triglyceride(s)
 - postprandial, *see* Postprandial triglycerides
 - relationship to soluble E-selectin, leptin, and insulin resistance, in nonobese Japanese with NIDDM, 376–380
 - response to high MUFA and high CHO diet, correlation to variations in plasma apo C-III levels, 1390–1397
 - sTNF-R, BMI, leptin and adiponectin in nonobese Japanese NIDDM patients, 1297–1301
 - and VLDL-C ratio, impact of apo E genotype, 1017
- Tumor growth factor β 1 (TGF- β 1), release by human AT, effect of obesity, 1546–1551
- Tumor necrosis factor- α (TNF- α)
- activity, relation to insulin resistance in nonobese Japanese NIDDM patients, 1297–1301
 - and high-sensitive CRP and cardiovascular risk factors before and after weight loss in obese children, 1155–1161
 - leptin expression associated with, in dimethylnitrosamine and bile duct ligation-induced liver cirrhosis, 445–452
 - reciprocal association with insulin resistance and C-reactive protein, in clinically healthy men, 439–444

- Turkish subjects, prepubertal obese children, endothelial activation and inflammation in, 1384-1389
- TVP, *see* Texturized vegetable protein
- Type 2 diabetes mellitus, *see* Non-insulin-dependent diabetes mellitus
- TZDs, *see* Thiazolidinedione(s)
- UC, *see* Unesterified cholesterol
- UCP3, *see* Uncoupling protein 3
- Ultrasonography, for evaluation of visceral fat and MS, 1230-1235
- Uncoupling protein 3 (UCP3), expression in skeletal muscle, and fat accumulation, effect of NO-1886 on HF diet-induced obesity, 1587-1592
- Unesterified cholesterol (UC) efflux, effects of HDL APF, 1087-1094
- Unipolar depression, *see* Major depressive disorder
- α,β (alpha,beta)-unsaturated aldehyde, 4-hydroxy-2-nonenal (HNE), effect on 2,6-ST mRNA expression in CYP2E1 cells, 733
- Uric acid
 serum, *see* Serum uric acid
 urinary excretion and plasma concentrations, influence of allopurinol on urinary purine loss after repeated sprint exercise in man, 1271-1272
 see also Hyperuricemia
- Urinary excretion
 of carnitine, glycine betaine and *N,N*-dimethylglycine, in patients attending a lipid clinic, 1-14
 of HGA in alkaptonuria patients, effect of nitisinone, 719-728
 of inosine, Hx and uric acid, influence of allopurinol on urinary purine loss after repeated sprint exercise in man, 1272-1273
 see also Albuminuria; Alkaptonuria; Micoalbuminuria
- Urinary isoprostane, association with adiponectin or leptin in Japanese Americans, 194-199
- Urinary purine loss, after repeated sprint exercise in man, influence of allopurinol, 1269-1275
- Vagal nerve stimulation, modifying effect on postprandial insulin sensitivity, 579-583
- Vascular adhesion molecule1 (VCAM-1), *see* Soluble vascular adhesion molecule1
- Vascular compliance, effect of HF meal in subgroup of young healthy subjects, 1337-1344
- Vascular endothelial growth factor (VEGF), release by human adipocytes, effect of insulin on, 220-226
- Vascular reactivity
 effects of FAs in healthy adult rats, 1600-1609
 effects of red wine consumption on, in NIDDM, 306-313
- Vascular smooth muscle proliferation, and serum Hcy and intimal hyperplasia, effect of rosiglitazone in high methionine-fed Sprague-Dawley rats, 645-652
- VAT, *see* Visceral adipose tissue
- VEGF, *see* Vascular endothelial growth factor
- Ventricular fibrillation (VF), outcomes in animal models, systematic review and meta-analysis of impact of ω -3 fatty acids, 1557-1565
- Ventricular tachycardia (VT), outcomes in animal models, systematic review and meta-analysis of impact of ω -3 fatty acids, 1557-1565
- Very long chain fatty acids (VLCFAs) (policosanols), and phytosterols, effects on plasma lipid levels and cholesterol biosynthesis, 508-814
- Very low calorie diet (VLCD), and withdrawal of blood glucose-lowering therapy, effect on endogenous glucose production in obese NIDDM, 705-712
- Very-low-density lipoprotein-cholesterol (VLDL-C), and TG ratio, impact of apo E genotype, 1017
- Very low-energy diet (VLED), 2-day, effect on SM insulin sensitivity in obese NIDDM patients on insulin therapy, 1669-1678
- Visceral adipose tissue (VAT)
 accumulation, and UCP3 expression in skeletal muscle, effect of NO-1886 on HF diet-induced obesity, 1587-1592
 effect of TZDs on, 24-32
 and MS, ultrasonography for evaluation of, 1230-1235
- VLCD, *see* Very low calorie diet
- VLCFAs, *see* Very long chain fatty acids
- VLDL-C, *see* Very-low-density lipoprotein-cholesterol
- VLED, *see* Very low-energy diet
- Voglibose, and acarbose, effects on postprandial hyperglycemia and subjective abdominal symptoms, 387-390
- Waist circumference (WC)
 and birth weight, and serum hepatic enzyme levels, of 18-year old sons, associations with their father's and mother's BMI, 466-470
 and BMI
 elderly versus young men with similar and smaller, oxidative and nonoxidative glucose disposal in, 748-755
 relationship between IMGU and excess adiposity, 1323-1328
- Waist-to-hip ratio (WHR), association with circulating OPG in healthy Korean women, 49-54
- Walking, fuel oxidation during, 793-799
- Walking time, for high serum uric acid levels, and alcohol drinking, possible interactions with eNOS genotype among Japanese, 1302-1308
- WAT, *see* White adipose tissue
- WBC, *see* White blood cell
- WC, *see* Waist circumference
- Weight gain, in NIDDM patients, role of insulinemia and sulfonylurea usage in, 28-29
- Weight loss
 associated with reduced intake of CHO, effect on atherogenicity of LDL in premenopausal women, 1133-1141

Weight loss (continued)

- high-sensitive CRP, TNF- α and cardiovascular risk factors in obese children, 1155-1161
- modest, effect on insulin action in obese African Americans, 960-965

Westernized lifestyle, effect on the association between fasting serum nonesterified fatty acids and insulin secretion in Japanese men, 713-718

White adipose tissue (WAT)

- of aging rats, effect of leptin on lipogenic enzyme expression through modification of SREBP-1c gene expression in, 1041-1047
- role in neurophysiologic modulation of obesity, 1204

White blood cell (WBC) count, normal, relation to cardiorespiratory fitness in apparently healthy Korean men, 1448-1452

Whole-blood viscosity, at multiple shear rates, influence of Lp, 764-768

Whole-body glucose metabolism, and insulin signaling, effect of ET antagonism in Zucker fatty rats, 1515-1523

Whole-body insulin sensitivity (S_i)

- of male offspring of high-fat fed rats, 500-507
- and muscle glucose transport, effect of selective angiotensin II receptor antagonism on hypertensive TG(mREN2)27 rats, 1659-1668

Whole-body protein anabolism, in response to insulin, influence of sex, 1529-1535

Whole-body protein turnover, effects of GH and/or sex steroids in healthy aged men and women, 1162-1167

WHR, *see* Waist-to-hip ratio

Women

- with and without GDM, MS at follow-up in index pregnancy, 1115-1121
- Korean, association of circulating OPG with age, WHR, serum TC, and LDL-C levels in, 49-54
- lean versus obese, ketone body metabolism in, 1542-1545
- long-term effect of recombinant methionyl human leptin therapy on hyperandrogenism and menstrual function and pituitary functions, 255-263

Women (continued)

- and men, healthy aged, effects of GH and/or sex steroids on whole-body protein turnover, 1162-1167
- with NIDDM, effect of single versus multiple bouts of exercise, 989-994
- obese and lean African American and white American, eNOS content in AT, 1368-1373
- with PCOS, and IR, metformin and diet benefits in, 113-121
- postmenopausal, *see* Postmenopausal (PM) women
- premenopausal, effect of weight loss associated with reduced intake of CHO on atherogenicity of LDL in, 1133-1141
- reproductive mood disorders in, S1:49-52
- see also* Sex

Xanthomatosis, familial massive tendon, with decreased HDL-mediated cholesterol efflux, 1095-1101

Xylitol, and ethanol, effect of simultaneous dietary administration on bone resorption, 548-551

Young subjects

men

- versus elderly with similar and smaller BMI and waist circumferences, oxidative and nonoxidative glucose disposal in, 748-755
- Japanese, cellular insulin resistance in EBV-transformed lymphoblasts obtained from, 370-375
- with LBW, susceptibility to diabetogenic effects of prolonged FFA exposure, 1398-1406
- and middle-aged athletes versus sedentary men, effects of exercise at moderate and hard intensity on substrate oxidation, 1411-1419
- waist circumference, birth weight and serum hepatic enzyme levels of, associations with their father's and mother's BMI, 466-470
- with pituitary gigantism, cardiac and metabolic effects of chronic GH and IGF-1 excess in, 1174-1180

Z-DNA-forming motif of promoter of *SLC11A1* gene, functional polymorphism of, association with IDDM in Japanese, 628-633

